

CANONICAL BASES OF SINGULARITY RINGEL-HALL ALGEBRAS AND HALL POLYNOMIALS

GUANGLIAN ZHANG

ABSTRACT. In this paper, the singularity Ringel-Hall algebras are defined. A new class of perverse sheaves are shown to have purity property. The canonical bases of singularity Ringel-Hall algebras are constructed. As an application, the existence of Hall polynomials in the tame quiver algebras is proved.

0. Introduction

0.1 Since G. Lusztig [L1, L2, L3] and M. Kashiwara [K] have proved that there exist the canonical bases in the quantized universal enveloping algebras, the canonical bases is playing an extremely vital role in the research of Lie algebra, Heck algebra and the quantized Schur algebra. The elements in the canonical bases are not only characteristic as perverse sheaves of quiver variety [L1, L2, L3], but also they are characteristic as the elements in Ringel-Hall algebra (see [L4] and [LXZ]).

Because of the deep relations between Ringel-Hall algebra and the many profound results in the representation theory of finite dimensional algebra, it is worth to study the representations of quiver algebras using the canonical bases and study the canonical bases and the representation of the other algebras (e.g., q -Schur algebra) using the representations of quivers.

M. Varagnolo and E. Vasserot [VV] proved the decomposition conjecture for quantized Schur algebra of type A .

A natural question is that how one can prove the decomposition conjecture for quantized Schur algebra of the other types.

Using the Ringel-Hall algebra model in the study of canonical bases is an important skill in [VV]. Let $\mathbf{U}_n^+(Q)$ (resp. $\mathbf{U}_\infty^+(Q)$) be the generic Hall algebra, where $Q = \tilde{A}_n$ (resp. $Q = A_\infty$). Based on the proof in [VV], we had to construct q -Fock spaces of the other affine types in order to prove the decomposition conjecture for quantized Schur algebra of the general type.

According to Proposition 5 in [VV], we know that $\mathbf{U}_\infty^+(Q)/I \cong \bigwedge^\infty$, where I is the submodule of $\mathbf{U}_\infty^+(Q)$ generated by $\mathbf{b}_\mathcal{O}$, where the $\mathcal{O}$'s are unstable orbits in Nakajima variety of type A_∞ . We conjecture that the q -Fock spaces of the other affine types should be the quotient of some subalgebra of Ringel-Hall algebra (e.g., Singularity Ringel-Hall algebras) modulo unstable orbits. In this paper, we prove that the quotient of the singularity Ringel-Hall algebras modulo unstable orbits is isomorphic to q -Fock space in the case of type A .

G.Lusztig [L4] proved that the perverse sheaves corresponding to the canonical bases have purity property. In this paper, we find a new class of perverse sheaves which has purity property.

The research was supported in part by NSF grant 10771112.

The research was also supported in part by Research Institute for Mathematical Sciences, Kyoto University, Kyoto and Chern Institute of Mathematics, Nankai University, Tianjin, China.

Thereby, we prove that the singularity Ringel-Hall algebras have the canonical bases. As an application of the singularity Ringel-Hall algebras, we prove that the tame quiver algebras have Hall polynomials.

0.2 The paper is organized as follows. In §1 we give a quick review of the definitions of Ringel-Hall algebras and Double Ringel-Hall algebras. In §2 we define the singularity Ringel-Hall algebras $\mathcal{H}^s(\Lambda)$ and study the relations between the singularity Ringel-Hall algebras and the quantized universal enveloping algebras. We prove Proposition 2.1.2 and from this, we point out that the quotient of the singularity Ringel-Hall algebras modulo unstable orbits is isomorphic to q -Fock space in the case of type A . In §3 we construct the PBW -basis of $\mathcal{H}^s(\Lambda)$. In §4 we prove that the closure of semi-simple objects in $\mathcal{T}_i$ have purity property. In §5 we study the fibres of p_3 . We also give a new class of perverse sheaves which have purity property. In §6 we prove that the singularity Ringel-Hall algebras have the canonical bases. In §7 A. Hubery [H] have proved the existence of Hall polynomials on the tame quivers for Segre classes. In this subsection, by using the extension algebras of singularity Ringel-Hall algebras, we give a simple and direct proof for the existence of Hall polynomials on the tame quivers.

Acknowledgments. I would like to express my sincere gratitude to H. Nakajima for a number of interesting discussions. I am also grateful to J. Xiao for a number of interesting discussions.

1. Ringel-Hall algebras

1.1 A quiver $Q = (I, H, s, t)$ consists of a vertex set I , an arrow set H , and two maps $s, t : H \rightarrow I$ such that an arrow $\rho \in H$ starts at $s(\rho)$ and terminates at $t(\rho)$.

Throughout the paper, $\mathbb{F}_q$ denotes a finite field with q elements, $k = \overline{\mathbb{F}}_q$ the algebraic closed field, $Q = (I, H, s, t)$ is a fixed connected quiver, and $\Lambda = \mathbb{F}_q Q$ is the path algebra of Q over $\mathbb{F}_q$. By $\text{mod } \Lambda$ we denote the category of all finite dimensional left Λ -modules, or equivalently finite modules. It is well-known that $\text{mod } \Lambda$ is equivalent to the category of finite dimensional representations of Q over $\mathbb{F}_q$. We shall simply identify Λ -modules with representations of Q .

The set of isomorphism classes of (nilpotent) simple Λ -modules is naturally indexed by the set I of vertices of Q . Then the Grothendieck group $G(\Lambda)$ of $\text{mod } \Lambda$ is the free Abelian group $\mathbb{Z}I$. For each nilpotent Λ -module M , the dimension vector $\underline{\dim} M = \sum_{i \in I} (\dim M_i) i$ is an element of $G(\Lambda)$. The Ringel-Hall algebra $\mathcal{H}(\Lambda)$ is graded by $\mathbb{N}I$, more precisely, by dimension vectors of modules.

The Euler form $\langle -, - \rangle$ on $G(\Lambda) = \mathbb{Z}I$ is defined by

$$\langle \alpha, \beta \rangle = \sum_{i \in I} a_i b_i - \sum_{\rho \in H} a_{s(\rho)} b_{t(\rho)}$$

for $\alpha = \sum_{i \in I} a_i i$ and $\beta = \sum_{i \in I} b_i i$ in $\mathbb{Z}I$. For any nilpotent Λ -modules M and N one has

$$\langle \underline{\dim} M, \underline{\dim} N \rangle = \dim_{\mathbb{F}_q} \text{Hom}_{\Lambda}(M, N) - \dim_{\mathbb{F}_q} \text{Ext}_{\Lambda}(M, N).$$

The symmetric Euler form is defined as

$$(\alpha, \beta) = \langle \alpha, \beta \rangle + \langle \beta, \alpha \rangle \quad \text{for } \alpha, \beta \in \mathbb{Z}I.$$

This gives rise to a symmetric generalized Cartan matrix $C = (a_{ij})_{i, j \in I}$ with $a_{ij} = (i, j)$. It is easy to see that C is independent of the field $\mathbb{F}_q$ and the orientation of Q .

Throughout the paper, we concentrate on tame quiver Q . The symmetric Euler forms give rise to the Cartan matrices of types A, D and E .

1.2 Ringel-Hall algebra. Given three modules L, M, N in $\text{mod } \Lambda$, let g_{MN}^L denote the number of Λ -submodules W of L such that $W \simeq N$ and $L/W \simeq M$ in $\text{mod } \Lambda$. More generally, for $M_1, \dots, M_t, L \in \text{mod } \Lambda$, let $g_{M_1 \dots M_t}^L$ denote the number of the filtrations $0 = L_0 \subseteq L_1 \subseteq \dots \subseteq L_t = L$ of Λ -submodules such that $L_i/L_{i-1} \simeq M_i$ for $i = 1, \dots, t$. Let $v_q = \sqrt{q} \in \mathbb{C}$ and $\mathcal{P}$ be the set of isomorphism classes of finite dimensional nilpotent Λ -modules. Then the twisted Ringel-Hall algebra $\mathcal{H}^*(\Lambda)$ is defined by setting $\mathcal{H}^*(\Lambda) = \mathcal{H}(\Lambda)$ as $\mathbb{Q}(v_q)$ -vector space, but the multiplication is defined by

$$u_{[M]} * u_{[N]} = v_q^{\langle \underline{\dim} M, \underline{\dim} N \rangle} \sum_{[L] \in \mathcal{P}} g_{MN}^L u_L.$$

Following [R3], for any Λ -module M , we denote $\langle M \rangle = v^{-\dim M + \dim \text{End}_\Lambda(M)} u_{[M]}$. Note that $\{\langle M \rangle \mid M \in \mathcal{P}\}$ a $\mathbb{Q}(v_q)$ -basis of $\mathcal{H}^*(\Lambda)$.

The $\mathbb{Q}(v_q)$ -algebras $\mathcal{H}^*(\Lambda)$ depends on q . We will use $\mathcal{H}_q^*(\Lambda)$ to indicate the dependence on q when such a need arises.

1.3 A construction by Lusztig. For any finite dimensional I -graded k -vector space $V = \sum_{i \in I} V_i$ with a given $\mathbb{F}_q$ -rational structure by Frobenius map F , let $\mathbb{E}_V$ be the subset of $\oplus_{\rho \in H} \text{Hom}(V_{s(\rho)}, V_{t(\rho)})$ defining nilpotent representations of Q . Note that $\mathbb{E}_V = \oplus_{\rho \in H} \text{Hom}(V_{s(\rho)}, V_{t(\rho)})$ when Q has no oriented cycles. The space of $\mathbb{F}_q$ -rational points of $\mathbb{E}_V$ is the fixed point set $\mathbb{E}_V^F$.

Let $G_V = \prod_{i \in I} GL(V_i)$, its subgroup of $\mathbb{F}_q$ -rational points is G_V^F . Then the group $G_V = \prod_{i \in I} GL(V_i)$ acts naturally on $\mathbb{E}_V$ by

$$(g, x) \mapsto g \bullet x = x' \quad \text{where} \quad x'_\rho = g_{t(\rho)} x_\rho g_{s(\rho)}^{-1} \quad \text{for all } \rho \in H.$$

The restriction of the action of G_V on $\mathbb{E}_V$ gives an action of the finite group G_V^F on $\mathbb{E}_V^F$.

For $\gamma \in \mathbb{N}I$, we fix a I -graded k -vector space V_γ with $\underline{\dim} V_\gamma = \gamma$. We set $\mathbb{E}_\gamma = \mathbb{E}_{V_\gamma}$ and $G_\gamma = G_{V_\gamma}$. For $\alpha, \beta \in \mathbb{N}I$ and $\gamma = \alpha + \beta$, we consider the diagram

$$\mathbb{E}_\alpha \times \mathbb{E}_\beta \xleftarrow{p_1} \mathbb{E}' \xrightarrow{p_2} \mathbb{E}'' \xrightarrow{p_3} \mathbb{E}_\gamma.$$

Here $\mathbb{E}''$ is the set of all pairs (x, W) , consisting of $x \in \mathbb{E}_\gamma$ and an x -stable I -graded subspace W of V_γ with $\underline{\dim} W = \beta$, and $\mathbb{E}'$ is the set of all quadruples (x, W, R', R'') , consisting of $(x, W) \in \mathbb{E}''$ and two invertible linear maps $R' : \mathbb{F}_q^\beta \rightarrow W$ and $R'' : \mathbb{F}_q^\alpha \rightarrow \mathbb{F}_q^\gamma/W$. The maps are defined in an obvious way: $p_2(x, W, R', R'') = (x, W)$, $p_3(x, W) = x$, and $p_1(x, W, R', R'') = (x', x'')$, where $x_\rho R'_{s(\rho)} = R'_{t(\rho)} x'_\rho$ and $x_\rho R''_{s(\rho)} = R''_{t(\rho)} x''_\rho$ for all $\rho \in H$.

If $M \in \mathbb{E}_\alpha, N \in \mathbb{E}_\beta$ and $L \in \mathbb{E}_{\alpha+\beta}$, we define

$$\mathbf{Z} = p_2 p_1^{-1}(\mathcal{O}_M \times \mathcal{O}_N), \mathbf{Z}_{L,M,N} = \mathbf{Z} \cap p_3^{-1}.$$

The varieties and morphisms in above diagram are naturally defined over $\mathbb{F}_q$. So we have

$$\mathbb{E}_\alpha^F \times \mathbb{E}_\beta^F \xleftarrow{p_1} \mathbb{E}'^F \xrightarrow{p_2} \mathbb{E}''^F \xrightarrow{p_3} \mathbb{E}_\gamma^F.$$

For any map $p : X \rightarrow Y$ of finite sets, $p^* : \mathbb{C}(Y) \rightarrow \mathbb{C}(X)$ is defined by $p^*(f)(x) = f(p(x))$ and $p_! : \mathbb{C}(X) \rightarrow \mathbb{C}(Y)$ is defined by $p_!(h)(y) = \sum_{x \in p^{-1}(y)} h(x)$, on the integration along the fibers. Let $\mathbb{C}_{G^F}(\mathbb{E}_V^F)$ be the space of G_V^F -invariant functions $\mathbb{E}_V^F \rightarrow \mathbb{C}$ (or $\overline{\mathbb{Q}_l}$.) Given $f \in \mathbb{C}_{G^F}(\mathbb{E}_\alpha^F)$ and $g \in \mathbb{C}_{G^F}(\mathbb{E}_\beta^F)$, there is a unique $h \in \mathbb{C}_{G^F}(\mathbb{E}''^F)$ such that $p_2^*(h) = p_1^*(f \times g)$. Then define $f \circ g$ by

$$f \circ g = (p_3)_!(h) \in \mathbb{C}_{G^F}(\mathbb{E}_\gamma^F).$$

Let

$$\mathbf{m}(\alpha, \beta) = \sum_{i \in I} a_i b_i + \sum_{\rho \in H} a_{s(\rho)} b_{t(\rho)}.$$

We again define the multiplication in the $\mathbb{C}$ -space $\mathbf{K} = \oplus_{\alpha \in \mathbb{N}I} \mathbb{C}_{GF}(\mathbb{E}_\alpha^F)$ by

$$f * g = v_q^{-\mathbf{m}(\alpha, \beta)} f \circ g$$

for all $f \in \mathbb{C}_{GF}(\mathbb{E}_\alpha^F)$ and $g \in \mathbb{C}_{GF}(\mathbb{E}_\beta^F)$. Then $(\mathbf{K}, *)$ becomes an associative $\mathbb{C}$ -algebra.

For $M \in \mathbb{E}_\alpha^F$, let $\mathcal{O}_M \subset \mathbb{E}_\alpha$ be the G_α -orbit of M . We take $\mathbf{1}_{[M]} \in \mathbb{C}_{GF}(\mathbb{E}_\alpha^F)$ to be the characteristic function of $\mathcal{O}_M^F$, and set $f_{[M]} = v_q^{-\dim \mathcal{O}_M} \mathbf{1}_{[M]}$. We consider the subalgebra $(\mathbf{L}, *)$ of $(\mathbf{K}, *)$ generated by $f_{[M]}$ over $\mathbb{Q}(v_q)$, for all $M \in \mathbb{E}_\alpha^F$ and all $\alpha \in \mathbb{N}I$. In fact $\mathbf{L}$ has a $\mathbb{Q}(v_q)$ -basis $\{f_{[M]} | M \in \mathbb{E}_\alpha^F, \alpha \in \mathbb{N}I\}$. Since $\mathbf{1}_{[M]} \circ \mathbf{1}_{[N]}(W) = g_{MN}^W$ for any $W \in \mathbb{E}_\gamma^F$, we have

Proposition 1.3.1 [LXZ] *The linear map $\varphi : (\mathbf{L}, *) \longrightarrow \mathcal{H}^*(\Lambda)$ defined by*

$$\varphi(f_{[M]}) = \langle M \rangle, \quad \text{for all } [M] \in \mathcal{P}$$

is an isomorphism of the associative $\mathbb{Q}(v_q)$ -algebras.

1.4 Double Ringel-Hall algebra $\mathcal{D}(\Lambda)$. First, we define a Hopf algebra $\mathcal{H}^+(\Lambda)$ which is a $\mathbb{Q}(v)$ -vector space with the basis $\{K_\mu | \mu \in \mathbb{Z}[I], \alpha \in \mathcal{P}\}$, whose Hopf algebra structure is given as

(a) Multiplication ([R1])

$$\begin{aligned} u_\alpha^+ * u_\beta^+ &= v^{\langle \alpha, \beta \rangle} \sum_{\lambda \in \mathcal{P}} g_{\alpha\beta}^\lambda u_\lambda^+, \quad \text{for all } \alpha, \beta \in \mathcal{P}, \\ K_\mu * u_\alpha^+ &= v^{(\mu, \alpha)} u_\alpha^+ * K_\mu, \quad \text{for all } \alpha \in \mathcal{P}, \mu \in \mathbb{N}[I], \\ K_\mu * K_\nu &= K_\nu * K_\mu = K_{\mu+\nu}, \quad \text{for all } \mu, \nu \in \mathbb{N}[I]. \end{aligned}$$

(b) Comultiplication ([G])

$$\begin{aligned} \Delta(u_\lambda^+) &= \sum_{\alpha, \beta \in \mathcal{P}} v^{\langle \alpha, \beta \rangle} \frac{a_\alpha a_\beta}{a_\lambda} g_{\alpha\beta}^\lambda u_\alpha^+ K_\beta \otimes u_\beta^+, \quad \text{for all } \lambda \in \mathcal{P}, \\ \Delta(K_\mu) &= K_\mu \otimes K_\mu, \quad \text{for all } \mu \in \mathbb{N}[I]. \end{aligned}$$

with counite $\epsilon(u_\lambda^+) = 0$, for all $\lambda \neq 0 \in \mathcal{P}$, and $\epsilon(K_\mu) = 1$. Here a_λ denotes the cardinality of finite set $\text{Aut}_\Lambda(M)$ with $\dim M = \lambda$.

(c) Antipode ([X])

$$\begin{aligned} S(u_\lambda^+) &= \delta_{\lambda 0} + \sum (-1)^m \sum_{\pi \in \mathcal{P}, \lambda_1, \dots, \lambda_m \in \mathcal{P} \setminus \{0\}} \times \\ &\quad v^{2 \sum_{i < j} \langle \lambda_i, \lambda_j \rangle} \frac{a_{\lambda_1} \cdots a_{\lambda_m}}{a_\lambda} g_{\lambda_1 \cdots \lambda_m}^\pi g_{\lambda_1 \cdots \lambda_m}^\pi K_{-\lambda} u_\pi^+. \end{aligned}$$

for all $\lambda \in \mathcal{P}$, and $S(K_\mu) = K_{-\mu}$ for all $\mu \in \mathbb{Z}[I]$. It has the subalgebra generated by $\{u_\lambda | \lambda \in \mathcal{P}\}$, which is isomorphic to $\mathcal{H}^*(\Lambda)$.

Dually, we can define a Hopf algebra $\mathcal{H}^-(\Lambda)$. Following Ringel, we have a bilinear form $\varphi : \mathcal{H}^+(\Lambda) \times \mathcal{H}^-(\Lambda) \longrightarrow \mathbb{Q}(v)$ defined by

$$\varphi(K_\mu u_\alpha^+, K_\nu u_\beta^-) = v^{-(\mu, \nu) - (\alpha, \nu) + (\mu, \beta)} \frac{|V_\alpha|}{a_\alpha} \delta_{\alpha\beta}$$

for all $\mu, \nu \in \mathbb{Z}[I]$ and all $\alpha, \beta \in \mathcal{P}$. Thanks to [X], we can obtain the reduced Drinfeld double $\mathcal{D}(\Lambda)$ of Ringel-Hall algebra of Λ , and the triangular decomposition

$$\mathcal{D}(\Lambda) = \mathcal{H}^- \otimes \mathcal{T} \otimes \mathcal{H}^+,$$

where $\mathcal{T}$ denotes the torus subalgebra generated by $\{K_\mu : \mu \in \mathbb{Z}[I]\}$.

The subalgebra of $\mathcal{D}(\Lambda)$ generated by $\{u_i^\pm, K_\pm | i \in I\}$ is called the composition algebra of Λ denoted by $\mathcal{C}(\Lambda)$. It is also a Hopf algebra and admits a triangular decomposition

$$\mathcal{C}(\Lambda) = \mathcal{C}^-(\Lambda) \otimes \mathcal{T} \otimes \mathcal{C}^+(\Lambda),$$

where $\mathcal{C}^+(\Lambda)$ is the composition algebra generated by $u_i^+ : i \in I$, and $\mathcal{C}^-(\Lambda)$ is defined dually. Moreover, the restriction $\varphi : \mathcal{C}^+(\Lambda) \times \mathcal{C}^-(\Lambda) \rightarrow \mathbb{Q}(v)$ is non-degenerate (see [HX]).

In addition, $\mathcal{D}(\Lambda)$ also admits an involution ω defined by

$$\begin{aligned} \omega(u_\lambda^+) &= u_\lambda^-, \omega(u_\lambda^-) = u_\lambda^+, \text{ for all } \lambda \in \mathcal{P}; \\ \omega(K_\mu) &= K_{-\mu}, \text{ for all } \mu \in \mathbb{N}[I]. \end{aligned}$$

Then $\varphi(x, y) = (\omega(x), \omega(y))$. Obviously, ω induces an involution of $\mathcal{C}(\Lambda)$.

2. Singularity Ringel-Hall algebras

2.1 Singularity Ringel-Hall algebras $\mathcal{H}^s(\Lambda)$. Let $\{\mathcal{T}_1, \dots, \mathcal{T}_l\}$ be all nonhomogeneous tubes in $\text{mod-}\Lambda$ (in fact, $l \leq 3$), $r_i = r(\mathcal{T}_i)$ the period of $\mathcal{T}_i$. It is well known that $\sum_{i=1}^l (r_i - 1) = |I| - 2$. We define $\mathcal{H}^s(\Lambda)$ to be the subalgebra of $\mathcal{H}^*(\Lambda)$ generated by $\{u_i, u_{[M]} : i \in I, M \in \mathcal{T}_j, 1 \leq j \leq l\}$. It is called Singularity Ringel-Hall algebra of Λ . From the definition, it is clear that $\mathcal{H}^s(\Lambda)$ depends only on the type of the quiver Q , and does not depend on the finite field $\mathbb{F}_q$. We now set $\mathcal{D}^s(\Lambda)$ to be the subalgebra of $\mathcal{D}(\Lambda)$ generated by $\{u_i^\pm, u_{[M]}^\pm : i \in I, M \in \mathcal{T}_j, 1 \leq j \leq l\}$.

According to the AR -quiver of tame quiver (see [CB]), it is easy to see that:

Lemma 2.1.1 *Let $M \in \mathcal{T}_i$, for some $i, 1 \leq i \leq l$, and*

$$0 \longrightarrow M_2 \longrightarrow M \longrightarrow M_1 \longrightarrow 0$$

a short exact sequence. Then $M_1 \cong I_1 \oplus N_1$, $M \cong P_2 \oplus N_2$, where P_2 is preprojective, $N_1, N_2 \in \mathcal{T}_i$, and I_1 is preinjective. $\square$

Lemma 2.1.2 *$\mathcal{D}^s(\Lambda)$ is a Hopf algebra depends on the type of Q .*

Proof. By Lemma 2.1.1, it is easy to see that $\mathcal{D}^s(\Lambda)$ is closed under comultiplication. So it is sufficient to prove that $\mathcal{D}^s(\Lambda)$ is closed under antipode S .

Since comultiplication is an algebra homomorphism, it is sufficient to prove that $S(u_{[M]}^+) \in \mathcal{D}^s(\Lambda)$, for some $M \in \mathcal{T}_i, 1 \leq i \leq l$.

We may assume that $S(u_{[M']}^+) \in \mathcal{D}^s(\Lambda)$ for any $\underline{\dim} M' < \underline{\dim} M$. Because of $\mu(S \otimes 1)\Delta = \eta\epsilon$, we have

$$S(u_{[M]}^+) + S(K_{\underline{\dim} M})u_{[M]}^+ + \sum_{M_1, M_2 \neq 0} v^{\langle \underline{\dim} M_1, \underline{\dim} M_2 \rangle} \frac{a_{M_1} a_{M_2}}{a_M} g_{M_1 M_2}^M S(u_{[M_1]}^+) K_{\underline{\dim} M_2} u_{[M_2]}^+. \quad (2.1.1)$$

Suppose $M_1 \cong I_1 \oplus N_1, M \cong P_2 \oplus N_2$. Using induction on $\underline{\dim} M$, we have, by Lemma 2.1.1, $S(u_{[N_1]}^+) \in \mathcal{D}^s(\Lambda)$.

Since $u_{[N_1]}^+ * u_{[I_1]}^+ = v^{\langle \underline{\dim} N_1, \underline{\dim} I_1 \rangle} u_{M_1}^+$, and $S(u_{[I_1]}^+) \in \mathcal{D}^s(\Lambda)$ by Lemma 6.1 and 6.2 in [LXZ], we have $S(u_{[M_1]}^+) \in \mathcal{D}^s(\Lambda)$. Therefore (2.1.1) implies $S(u_{[M]}^+) \in \mathcal{D}^s(\Lambda)$. $\square$

2.2 Decomposition of $\mathcal{H}^s(\Lambda)$. In the following, we follow an idea of Sevenhant and Van den Bergh to obtain subalgebras of $\mathcal{H}^s(\Lambda)$ and $\mathcal{D}^s(\Lambda)$. (see also [HX].)

According to the definition of φ , it is easy to see that the restriction of φ on $\mathcal{H}_\alpha^{s,+} \times \mathcal{H}_\alpha^{s,-}$, for all $\alpha \in \mathbb{N}[I]$, is also non-degenerate.

For $\alpha, \beta \in \mathbb{N}[I]$, we use $\alpha \leq \beta$ to mean that $\beta - \alpha \in \mathbb{N}[I]$. Clearly, $\mathcal{C}(\Lambda)_\beta = \mathcal{H}^s(\Lambda)_\beta$ if $\beta < \delta$. We now define

$$\mathcal{L}_\delta^+ = \{x^+ \in \mathcal{H}^{s,+}(\Lambda) \mid \varphi(x^+, \mathcal{C}^-(\Lambda)) = 0\}.$$

Dually, let

$$\mathcal{L}_\delta^- = \{x^- \in \mathcal{H}^{s,-}(\Lambda) \mid \varphi(x^-, \mathcal{C}^+(\Lambda)) = 0\}.$$

The non-degeneracy of φ implies

$$\mathcal{H}^{s,+}(\Lambda)_\delta = \mathcal{C}^+(\Lambda)_\delta \oplus \mathcal{L}_\delta^+,$$

$$\mathcal{H}^{s,-}(\Lambda)_\delta = \mathcal{C}^-(\Lambda)_\delta \oplus \mathcal{L}_\delta^-.$$

Let $\mathcal{D}^s(1)$ be the subalgebra of $\mathcal{D}^s$ generated by $\mathcal{C}^\pm(\Lambda)$ and $\mathcal{L}_\delta^\pm$. Then we also have the triangular decomposition

$$\mathcal{D}^s(1) = \mathcal{D}^s(1)^- \otimes \mathcal{T} \otimes \mathcal{D}^s(1)^+.$$

Suppose $\mathcal{L}_{(m-1)\delta}^\pm$ has been defined, we inductively define $\mathcal{L}_{m\delta}^\pm$ as follows:

$$\mathcal{L}_{m\delta}^+ = \{x^+ \in \mathcal{H}_{m\delta}^{s,+} \mid \varphi(x^+, \mathcal{D}^s(m-1)^-) = 0\},$$

$$\mathcal{L}_{m\delta}^- = \{x^- \in \mathcal{H}_{m\delta}^{s,-} \mid \varphi(x^-, \mathcal{D}^s(m-1)^+) = 0\}.$$

Let $\mathcal{D}^s(m)$ be the subalgebra of $\mathcal{D}^s$ generated by $\mathcal{D}^s(m-1)^\pm$ and $\mathcal{L}_{m\delta}^\pm$. Therefore

$$\mathcal{D}^s(m) = \mathcal{D}^s(m)^- \otimes \mathcal{T} \otimes \mathcal{D}^s(m)^+.$$

Lemma 2.2.1 *Let $\eta_{n\delta} = \dim_{\mathbb{Q}(v)} \mathcal{L}_{n\delta}^+ = \dim_{\mathbb{Q}(v)} \mathcal{L}_{n\delta}^-$. Then $\eta_{n\delta} = l$.*

Proof. By proposition 7.5 in [LXZ], we know that $\Psi(u_{[M(\alpha)]})$ for $\alpha \in \Phi_{Prep}^+$; $\Psi(u_{\alpha,i})$ for $\alpha \in \mathcal{T}_i$, the real roots: $i = 1, \dots, l$; $\Psi(u_{j,m\delta,i} - u_{j+1,m\delta,i})$, $m \geq 1$, $1 \leq j \leq r_i - 1$, $i = 1, \dots, l$; $\Psi(\tilde{E}_{n\delta})$, $n \geq 1$ and $\Psi(u_{[M(\beta)]})$ for $\beta \in \Phi_{Prei}^+$ form a $\mathbb{Z}$ -basis of $\mathfrak{n}^+$, that is, the basis of $\mathcal{C}^+(\Lambda)/(v-1)$.

According to the definition of $\mathcal{H}^s$, $\Psi(u_{[M(\alpha)]})$ for $\alpha \in \Phi_{Prep}^+$; $\Psi(u_{\alpha,i})$ for $\alpha \in \mathcal{T}_i$ the real root: $i = 1, \dots, l$; $\Psi(u_{j,m\delta,i})$, $m \geq 1$, $1 \leq j \leq r_i$, $i = 1, \dots, l$; $\Psi(\tilde{E}_{n\delta})$, $n \geq 1$ and $\Psi(u_{[M(\beta)]})$ for $\beta \in \Phi_{Prei}^+$ form a $\mathbb{Z}$ -basis of $\mathcal{H}^s/(v-1)$.

Since $\sum_{i=1}^l (r_i - 1) = |I| - 2$, we have $\eta_{n\delta} = l$ by the construction of $\mathcal{L}_{n\delta}^\pm$ for all n . $\square$

For each $n\delta$, there exist bases $\{x_n^1, \dots, x_n^l\}$ of $\mathcal{L}_{n\delta}^+$ and a basis $\{y_n^1, \dots, y_n^l\}$ of $\mathcal{L}_{n\delta}^-$ such that

$$\varphi(x_n^p, y_n^q) = \frac{1}{v - v^{-1}} \delta_{pq}.$$

Thus, we have

$$x_n^p y_m^q - y_m^q x_n^p = \frac{K_{n\delta} - K_{-n\delta}}{v - v^{-1}} \delta_{pq} \delta_{mn},$$

for all $m, n, 1 \leq p, q \leq l$ (see [HX]).

We now set $x_i = u_i^+$, $y_i = -v^{-1} u_i^-$ for each $i \in I$, and let $J = \{(n\delta, p) : 1 \leq p \leq l\}$. Define

$$\theta_i = i \text{ if } i \in I, \theta_i = n\delta \text{ if } i \in J, \text{ and } x_i = x_n^p \text{ if } i = (n\delta, p), y_i = -v^{-1} y_n^p \text{ if } i = (n\delta, p).$$

Moreover, by a theorem of Sevenhant and Van den Bergh, we have that $\mathcal{D}^s$ is generated by

$$\{x_i, y_i \mid i \in I \cup J\} \cup \{K_\mu : \mu \in \mathbb{N}[I]\}$$

with the defining relations

$$K_0 = 1, K_\mu K_\nu = K_{\mu+\nu} \text{ for all } \mu, \nu \in \mathbb{N}[I] \quad (2.2.1)$$

$$K_\mu x_i = v^{(\mu, \theta_i)} x_i K_\mu, \text{ and } K_\mu y_i = v^{-(\mu, \theta_i)} y_i K_\mu \text{ for all } i \in I \cup J, \mu \in \mathbb{N}[I] \quad (2.2.2)$$

$$x_i y_j - y_j x_i = \frac{K_{\theta_i} - K_{-\theta_i}}{v - v^{-1}} \delta_{ij} \text{ for all } i, j \in I \cup J \quad (2.2.3)$$

$$\sum_{p+p'=1-a_{ij}} (-1)^p x_i^{(p)} x_j x_i^{(p')} = 0 \text{ and } \sum_{p+p'=1-a_{ij}} (-1)^p y_i^{(p)} y_j y_i^{(p')} = 0 \quad (2.2.4)$$

$$x_i x_j = x_j x_i \text{ and } y_i y_j = y_j y_i \text{ for all } i, j \in I \cup J \text{ with } (\theta_i, \theta_i) = 0. \quad (2.2.5)$$

Applying the relations above, the next statement is clear.

Proposition 2.2.2

- (a) $\{x_n^i \mid n \in \mathbb{N}, 1 \leq i \leq l\}$ are central in $\mathcal{H}^{s,+}$. Dually, $\{y_n^i \mid n \in \mathbb{N}, 1 \leq i \leq l\}$ are central in $\mathcal{H}^{s,-}$.
- (b) $\mathcal{H}^{s,+} \cong \mathcal{C}^+ \otimes \mathbb{Q}(v)[x_n^i \mid n \in \mathbb{N}, 1 \leq i \leq l]$.
- (c) x_n^j commutes with $\mathcal{H}^{s,-}$, y_n^j commutes with $\mathcal{H}^{s,+}$.

□

In particular, we have $l = 1$ in case $Q = \tilde{A}_n, n \geq 3$, that is $\mathcal{H}^{s,+} \cong U_q(\tilde{sl}_n)^+ \otimes \mathbb{Q}(v)[x_1, \dots, x_n, \dots]$.

3. PBW-bases of Singularity Ringel-Hall algebras

3.1 It is known that there exists full subcategory $\mathfrak{C}(P, L)$ in $\text{mod } \Lambda$ (see [LXZ] 7.1). Moreover $\mathfrak{C}(P, L)$ is equivalent to the module category of the Kronecker quiver over $\mathbb{F}_q$. Thus it induces an exact embedding $F : \text{mod } K \hookrightarrow \text{mod } \Lambda$, where K is the path algebra of the Kronecker quiver over $\mathbb{F}_q$. We note here that the embedding functor F is essentially independent of the field $\mathbb{F}_q$. This gives rise to an injective homomorphism of algebras, still denoted by $F : \mathcal{H}^*(K) \hookrightarrow \mathcal{H}^*(\Lambda)$. In $\mathcal{H}^*(K)$ we have defined the element $E_{m\delta_K}$ for $m \geq 1$. Set $E_{m\delta} = F(E_{m\delta_K})$. Since $E_{m\delta_K} \in \mathcal{C}^*(K)$, and $\langle L \rangle, \langle P \rangle \in \mathcal{C}^*(\Lambda)$, so $E_{m\delta}$ is in $\mathcal{C}^*(\Lambda)$ and even in $\mathcal{C}^*(\Lambda)_{\mathcal{Z}}$. Let $\mathcal{K}$ be the subalgebra of $\mathcal{C}^*(\Lambda)$ generated by $E_{m\delta}$ for $m \in \mathbb{N}$, it is a polynomial ring on infinitely many variables $\{E_{m\delta} \mid m \geq 1\}$, and its integral form is the polynomial ring on variables $\{E_{m\delta} \mid m \geq 1\}$ over $\mathcal{Z}$.

We denote by $\mathfrak{C}_0$ (resp. $\mathfrak{C}_1$) the full subcategory of $\mathfrak{C}(P, L)$ consisting of the Λ -modules which belong to homogeneous (resp. non-homogeneous) tubes of $\text{mod } \Lambda$.

We now decompose $E_{n\delta}$ as follows

$$E_{n\delta} = E_{n\delta,1} + E_{n\delta,2} + E_{n\delta,3},$$

where

$$E_{n\delta,1} = v^{-n \dim S_1 - n \dim S_2} \sum_{[M], M \in \mathfrak{C}_1, \underline{\dim} M = n\delta} u_{[M]} \quad (3.1.1)$$

$$E_{n\delta,2} = v^{-n \dim S_1 - n \dim S_2} \sum_{\substack{[M], \underline{\dim} M = n\delta \\ M = M_1 \oplus M_2, 0 \neq M_1 \in \mathfrak{C}_1, 0 \neq M_2 \in \mathfrak{C}_0}} u_{[M]} \quad (3.1.2)$$

$$E_{n\delta,3} = v^{-n \dim S_1 - n \dim S_2} \sum_{[M], M \in \mathfrak{C}_0, \underline{\dim} M = n\delta} u_{[M]}. \quad (3.1.3)$$

Note that $\dim S_i = \dim_{\mathbb{F}_q} S_i, i = 1, 2$, but the values are independent of the choice of finite field $\mathbb{F}_q$. Let $\mathbf{w} = (w_1, \dots, w_t)$ be a partition of n , we then define

$$E_{\mathbf{w}\delta,3} = E_{w_1\delta,3} * \dots * E_{w_t\delta,3}.$$

Let $\mathbf{P}(n)$ be the set of all partitions of n , and $\langle N \rangle = v^{-\dim N + \dim \text{End}(N)} u_{[N]}$. Set

$$\mathbf{B} = \{\langle P \rangle * \langle M \rangle * E_{\mathbf{w}\delta,3} * \langle I \rangle \mid P \in \mathcal{P}_{\text{prep}}, M \in \oplus_{i=1}^l \mathcal{T}_i, I \in \mathcal{P}_{\text{prei}}, \mathbf{w} \in \mathbf{P}(n), n \in \mathbb{N}\}.$$

Then we have the following:

Theorem 3.1.1 *The set $\mathbf{B}$ is $\mathbb{Q}(v)$ -bases of $\mathcal{H}^{s,+}$.*

Proof. Let Π_i^a be the set of aperiodic r_i -tuples of partitions, for all $1 \leq i \leq l$. Set

$$\begin{aligned} \mathbf{B}^c &= \{ \langle P \rangle * E_{\pi_1} * \cdots * E_{\pi_l} * E_{\mathbf{w}\delta} * \langle I \rangle : P \in \mathcal{P}_{prep}, I \in \mathcal{P}_{prei}, \pi_i \in \Pi_i^a, \mathbf{w} \in \mathbf{P}(n), 1 \leq i \leq l, n \in \mathbb{N} \}, \\ \mathbf{B}' &= \{ x * f(x_n^j) : x \in \mathbf{B}^c, f(x_n^j) \in \mathbb{Q}(v)[x_n^j : 1 \leq j \leq l, n \in \mathbb{N}] \}. \end{aligned}$$

By Proposition 7.2 in [LXZ], we know that $\mathbf{B}^c$ is a $\mathbb{Q}(v)$ -basis of $\mathcal{C}^*(\Lambda)$. Following Proposition 2.2.2, we conclude that $\mathbf{B}'$ is $\mathbb{Q}(v)$ -basis of $\mathcal{H}^{s,+}$.

Using induction on n , it follows from (3.1.1), (3.1.2) and (3.1.3) that $E_{n\delta,3} \in \mathcal{H}^{s,+}$. Therefore $\mathbf{B} \subseteq \mathcal{H}^{s,+}$.

Because $\mathbf{B}$ is linear independent over $\mathbb{Q}(v)$, we will see that each

$$\langle P \rangle * E_{\pi_1} * \cdots * E_{\pi_l} * E_{\mathbf{w}\delta} * \langle I \rangle * f(x_n^j)$$

in $\mathbf{B}'$ may be represented by $\mathbf{B}$.

If M is a homogeneous module in some tubes, N a regular module in other tubes, then $\langle M \rangle * \langle N \rangle = \langle N \rangle * \langle M \rangle$. Thus, (3.1.1), (3.1.2) and (3.1.3) imply that $E_{\mathbf{w}\delta}$ is represented by $\mathbf{B}$. Furthermore, $\mathbf{B}^c$ may also be represented by $\mathbf{B}$. So it is sufficient to prove that x_n^j satisfy the property. By the definition of $\mathcal{H}^{s,+}$, we know that $\mathcal{H}^{s,+}$ is generated by $\mathbf{B}^c$ and $\{ \langle M \rangle : M \in \bigoplus_{i=1}^l \mathcal{T}_i \}$. Note that $M_1 \in \mathcal{T}_i$ or $M_1 = 0$ if $0 \rightarrow P \rightarrow P_1 \oplus M_1 \rightarrow M \rightarrow 0$ is a short exact sequence, where $P, P_1 \in \mathcal{P}_{prep}, M \in \mathcal{T}_i$ for some i . Similarly, we have $M_1 \in \mathcal{T}_i$ or $M_1 = 0$ if $0 \rightarrow M \rightarrow I_1 \oplus M_1 \rightarrow I \rightarrow 0$ is a short exact sequence, where $I, I_1 \in \mathcal{P}_{prei}, M \in \mathcal{T}_i$. It implies that every element in $\mathcal{H}^{s,+}$, in particular x_n^j , may be represented by $\mathbf{B}$.

4. Purity Properties of Perverse sheaves of closure of semi-simple objects in $\mathcal{T}_i$

4.1 We denote by $M(x), x \in \mathbb{E}_\alpha$, the Λ -module of dimension vector α corresponding to x . For subsets $\mathcal{A} \subset \mathbb{E}_\alpha$ and $\mathcal{B} \subset \mathbb{E}_\beta$, we define the extension set $\mathcal{A} \star \mathcal{B}$ of $\mathcal{A}$ by $\mathcal{B}$ to be

$$\begin{aligned} \mathcal{A} \star \mathcal{B} &= \{ z \in \mathbb{E}_{\alpha+\beta} \mid \text{there exists an exact sequence} \\ &\quad 0 \rightarrow M(x) \rightarrow M(z) \rightarrow M(y) \rightarrow 0 \text{ with } x \in \mathcal{B}, y \in \mathcal{A} \}. \end{aligned}$$

It follows from the definition that $\mathcal{A} \star \mathcal{B} = p_3 p_2 (p_1^{-1}(\mathcal{A} \times \mathcal{B}))$. Because p_1 is a locally trivial fibration (see Lemma 2.3 in [LXZ]), we then have $\overline{\mathcal{A} \star \mathcal{B}} \subseteq \overline{\mathcal{A}} \star \overline{\mathcal{B}}$. In particular, $\overline{\mathcal{O}}_M \star \overline{\mathcal{O}}_N = \overline{\mathcal{O}_{M \oplus N}}$ if $\text{Ext}(M, N) = 0$, i.e., $\mathcal{O}_{M \oplus N}$ is open and dense in $\overline{\mathcal{O}}_M \star \overline{\mathcal{O}}_N$.

Set $\text{codim } \mathcal{A} = \dim \mathbb{E}_\alpha - \dim \mathcal{A}$. We will need the following:

Lemma 4.1.1 [Re] *Given any $\alpha, \beta \in \text{NI}$, if $\mathcal{A} \subset \mathbb{E}_\alpha$ and $\mathcal{B} \subset \mathbb{E}_\beta$ are irreducible algebraic varieties and are stable under the action of G_α and G_β respectively, then $\mathcal{A} \star \mathcal{B}$ is irreducible and stable under the action of $G_{\alpha+\beta}$, too. Moreover,*

$$\text{codim } \mathcal{A} \star \mathcal{B} = \text{codim } \mathcal{A} + \text{codim } \mathcal{B} - \langle \beta, \alpha \rangle + r,$$

where $0 \leq r \leq \min\{\dim_k \text{Hom}(M(y), M(x)) \mid y \in \mathcal{B}, x \in \mathcal{A}\}$. □

Let $M(x), N(y)$ be modules corresponding to $x, y \in \mathbb{E}_\alpha$ respectively. We denote by $\mathcal{O}_x$ (or $\mathcal{O}_{M(x)}$) the G_α -orbits of x . We now introduce two orders in Λ -mod as follows:

- $N \leq_{deg} M$ if $\mathcal{O}_N \subseteq \overline{\mathcal{O}}_M$.
- $N \leq_{ext} M$ if there exists M_i, U_i, V_i and short exact sequence

$$0 \rightarrow U_i \rightarrow M_i \rightarrow V_i \rightarrow 0$$

such that $M = M_1, M_{i+1} = U_i \oplus V_i, 1 \leq i \leq p$, and $N = M_{p+1}$ for some natural number p .

It follows from G.Zwara [Z] that:

Proposition 4.1.2 *The orders $\leq_{deg}, \leq_{ext}$ are equivalent in $\Lambda - mod$.* $\square$

We denote by $\mathcal{P}_X$ the category of perverse sheaves on algebraic variety X . Let f be a locally closed embedding from X to Y . One has the intermediate extension functor

$$f_{!*} : \mathcal{P}_X \longrightarrow \mathcal{P}_Y, P \longmapsto Im\{ {}^p\mathcal{H}^0(f_!P) \longrightarrow {}^p\mathcal{H}^0(f_*P) \}.$$

In particular, suppose $\mathcal{L}$ be a local system on a nonsingular Zariski dense open subset $j : U \longrightarrow Y$ of the irreducible n -dimensional Y . Then $IC_Y(\mathcal{L}) := j_{!*}\mathcal{L}[n] \in \mathcal{P}_Y$.

Definition 4.1.3 *Let $\mathcal{K}$ be a Weil complex. Then $\mathcal{K}$ is said to have the purity property if on all stalks of the semisimplification $\mathcal{K}_x^{ss}(v/2)$ of cohomology sheaves the Frobenius F_x acts trivially.*

Lemma 4.1.4 *Let $p : X \longrightarrow Y$ be a smooth morphism of relative dimension $d, \dim Y = m$, a nonsingular Zariski dense open subset U of the irreducible n -dimensional Y and the diagram as follows is a cartesian square*

$$\begin{array}{ccc} U = p^{-1}(U_0) & \xrightarrow{j} & X \\ p|_{U_0} \downarrow & & p \downarrow \\ U & \xrightarrow{j_0} & Y. \end{array}$$

If $j_{0!*}\overline{\mathbb{Q}}_l[m]$ has the purity property, then $j_{!*}\overline{\mathbb{Q}}_l[d+m]$ has the purity property.

Proof. By the definition of j_* and $j_!$, we have a natural morphism

$$\varphi_Y : {}^p\mathcal{H}^0(j_{0!*}\overline{\mathbb{Q}}_l[m]) \longrightarrow {}^p\mathcal{H}^0(j_{0*}\overline{\mathbb{Q}}_l[m]).$$

It induces an intermediate extension functor

$$j_{0!*} : \mathcal{P}_U \longrightarrow \mathcal{P}_Y$$

such that $j_{0!*}\overline{\mathbb{Q}}_l[m] = Im\{ {}^p\mathcal{H}^0(j_{0!}\overline{\mathbb{Q}}_l[m]) \longrightarrow {}^p\mathcal{H}^0(j_{0*}\overline{\mathbb{Q}}_l[m]) \}$. Furthermore,

$$p^*[d] \circ \varphi_Y : p^*[d]({}^p\mathcal{H}^0(j_{0!}\overline{\mathbb{Q}}_l[m]) \longrightarrow p^*[d]({}^p\mathcal{H}^0(j_{0*}\overline{\mathbb{Q}}_l[m])).$$

Since $p : X \longrightarrow Y$ is a smooth of relative dimension d , $p^*[d] = p^![-d]$ are t -exact, we have

$$\varphi_X : {}^p\mathcal{H}^0(p^*j_{0!}\overline{\mathbb{Q}}_l[m+d]) \longrightarrow {}^p\mathcal{H}^0(p^*j_{0*}\overline{\mathbb{Q}}_l[m+d]),$$

$$\varphi_X : {}^p\mathcal{H}^0(j_!\overline{\mathbb{Q}}_l[m+d]) \longrightarrow {}^p\mathcal{H}^0(j_*\overline{\mathbb{Q}}_l[m+d]).$$

So, $p^*[d](j_{0!*}\overline{\mathbb{Q}}_l[m]) = j_{!*}\overline{\mathbb{Q}}_l[d+m]$. By the same argument, we get $p^*[d] \circ F = F \circ p^*[d]$. Since $j_{0!*}\overline{\mathbb{Q}}_l[m]$ has the purity property, the statement of Lemma is true. $\square$

Let $\mathcal{N}_{w_i}$ and $\mathcal{N}_{w_i,3}$ are, respectively, the union of orbits of regular modules of $\mathfrak{C}(P, L)$ and $\mathfrak{C}_0(P, L)$ with dimension vector $w_i\delta$. Set $\mathcal{N}_{\mathbf{w}} = \mathcal{N}_{w_1} \star \cdots \star \mathcal{N}_{w_t}$ and $\mathcal{N}_{\mathbf{w},3} = \mathcal{N}_{w_1,3} \star \cdots \star \mathcal{N}_{w_t,3}$.

For any $P \in \mathcal{P}_{prep}$, $M \in \oplus_{i=1}^l \mathcal{T}_i$, $I \in \mathcal{P}_{prei}$, $\pi_i \in \Pi_i^a$, $1 \leq i \leq l$, $\mathbf{w} \in \mathbf{P}(n)$, $n \in \mathbb{N}$, we define the varieties

$$\mathcal{O}_{P,\pi_1,\dots,\pi_l,\mathbf{w},I} = \mathcal{O}_P \star \mathcal{O}_{\pi_1} \cdots \star \mathcal{O}_{\pi_l} \star \mathcal{N}_{\mathbf{w}} \star \mathcal{O}_I, \quad \mathcal{O}_{P,M,\mathbf{w},I} = \mathcal{O}_P \star \mathcal{O}_M \star \mathcal{N}_{\mathbf{w},3} \star \mathcal{O}_I.$$

According to [L4] and [L5], we know that $IC_{\overline{\mathcal{O}}_{P,\pi_1,\dots,\pi_l,\mathbf{w},I}}(\mathbb{Q}_l)$ have purity property. In order to construct the canonical basis of $\mathcal{H}^s(\Lambda)$, we need to study the purity property of $\overline{\mathcal{O}}_{P,M,\mathbf{w},I}$.

Theorem 4.1.5 *Let $X = \overline{\mathcal{O}}_{P,M,\mathbf{w},I}$, for all $P \in \mathcal{P}_{prep}$, $M \in \oplus_{i=1}^l \mathcal{T}_i$, $I \in \mathcal{P}_{prei}$, $\mathbf{w} \in \mathbf{P}(n)$, $n \in \mathbb{N}$. Then $IC_X(\mathbb{Q}_l)$ has the purity property.*

In order to prove Theorem 4.1.5, we need to prove a number of Lemmas about purity properties of perverse sheaves of closure of semi-simple objects in $\mathcal{T}_i$.

4.2 For the tame quivers $\tilde{A}_n, \tilde{D}_n, \tilde{E}_6, \tilde{E}_7$ and $\tilde{E}_8$. The orientations are given (see blow)




Fig.4.2.1




Fig.4.2.2




Fig.4.2.3




Fig.4.2.4




Fig.4.2.5

We then have (or following [DR])

(4.2.1) the real regular simples of $\tilde{A}_n$ have period n , and have dimension vectors $(1, 0, 0, \dots, 0, 1), (0, 1, 0, \dots, 0, 0), \dots, (0, 0, 0, \dots, 1, 0)$.

(4.2.2) the real regular simples of $\tilde{D}_n$ with dimension vectors

$(1, 0, 1, 1, \dots, 1, 1, 0), (0, 1, 1, 1, \dots, 1, 0, 1)$ (with period 2)

$(1, 0, 1, 1, \dots, 1, 0, 1), (0, 1, 1, 1, \dots, 1, 1, 0)$ (with period 2)

$(1, 1, 1, 0, \dots, 0, 0, 0), (0, 0, 1, 1, \dots, 1, 1, 1), (0, 0, 0, 1, 0, \dots, 0, 0, 0), \dots, (0, 0, 0, 0, \dots, 0, 1, 0, 0)$ (with period $n-2$)

(4.2.3) the real regular simples of $\tilde{E}_6$ have period 2, 3, 3 respectively, and have dimension vectors, respectively,

$$(1, 1, 2, 1, 1, 1, 1), (0, 1, 1, 1, 0, 1, 0); (1, 1, 1, 1, 0, 0, 0), (0, 1, 1, 0, 0, 1, 1), (0, 0, 1, 1, 1, 1, 0);$$

$$(1, 1, 1, 0, 0, 1, 0), (0, 1, 1, 1, 1, 0, 0), (0, 0, 1, 1, 0, 1, 1).$$

(4.2.4) the real regular simples of $\tilde{E}_7$ have period 2, 3, 4 respectively, and have dimension vectors, respectively,

$$(1, 1, 2, 2, 1, 1, 0, 1), (0, 1, 1, 2, 2, 1, 1, 1); (1, 1, 1, 2, 1, 1, 1, 1), (0, 1, 1, 1, 1, 1, 0, 0), (0, 0, 1, 1, 1, 0, 0, 1);$$

$$(1, 1, 1, 1, 1, 0, 0, 0), (0, 1, 1, 1, 0, 0, 0, 1), (0, 0, 1, 1, 1, 1, 1, 0), (0, 0, 0, 1, 1, 1, 0, 1).$$

(4.2.5) the real regular simples of $\tilde{E}_8$ have period 2, 3, 5 respectively, and have dimension vectors, respectively,

$$(1, 2, 3, 2, 2, 1, 1, 0, 2), (1, 2, 3, 3, 2, 2, 1, 1, 1);$$

$$(1, 2, 2, 1, 1, 1, 0, 0, 1), (0, 1, 2, 2, 2, 1, 1, 1, 1), (1, 1, 2, 2, 1, 1, 1, 0, 1);$$

$$(1, 1, 1, 1, 1, 0, 0, 0, 0), (0, 1, 1, 1, 0, 0, 0, 0, 1), (1, 1, 2, 1, 1, 1, 1, 1, 1), (0, 1, 1, 1, 1, 1, 1, 0, 0),$$

$$(0, 0, 1, 1, 1, 1, 0, 0, 1).$$

In the following, the $m \times n$ matrix over k is denoted by $M_{mn}(k)$, the k -vector space generated by column vectors of matrix A is denoted by $C(A)$, the projective space of k -vector space V is denoted by $P(V)$.

Lemma 4.2.1 *Let $Q = \tilde{A}_n, n \geq 3$ as in Fig.4.2.1. Let $E_i, 1 \leq i \leq n$, be regular simple objects, and $X = \overline{\mathcal{O}}_{\oplus_{i=1}^n E_i}$. Then $IC_X(\overline{\mathbb{Q}}_l)$ has the purity property.*

Proof. Let V be the I -graded vector space $V = \oplus_{i=1}^{n+1} V_i$, and $V_i = k$. Set $\alpha = (1, 1, \dots, 1) \in \mathbb{N}[I]$, then

$$\mathbb{E}_\alpha = \mathbb{E}_V = \{x \mid x = (x_{12}, x_{23}, \dots, x_{n,n+1}, x_{1,n+1}) \in k^{n+1}\},$$

$$\mathcal{O}_{\oplus_{i=1}^n E_i} = \{x \mid x_{12} = x_{23} = \dots = x_{n,n+1} = 0, x_{1,n+1} \neq 0\},$$

$$X = \overline{\mathcal{O}}_{\oplus_{i=1}^n E_i} = \{x \mid x_{12} = x_{23} = \dots = x_{n,n+1} = 0\} = \mathbb{A}^1.$$

It is clear that $IC_X(\mathbb{Q}_l)$ has the purity property. The statement is proved. $\square$

Lemma 4.2.2 *Let $Q = \tilde{D}_n, n \geq 4$ as in Fig.4.2.2, let $E_i, i = 1, 2; E'_j, j = 1, 2; E''_m, 1 \leq m \leq n-2$ be regular simple objects corresponding to three non-homogeneous tubes, respectively. Set $X_1 = \overline{\mathcal{O}}_{E_1 \oplus E_2}, X_2 = \overline{\mathcal{O}}_{E'_1 \oplus E'_2}$, and $X_3 = \overline{\mathcal{O}}_{\oplus_{i=1}^{n-2} E''_i}$. Then $IC_{X_i}(\overline{\mathbb{Q}}_l)$ have the purity property for any $i = 1, 2, 3$.*

Proof. We only prove $i = 1$. The other cases can be proved similarly. Let V be the I -graded vector space $V = \oplus_{i=1}^{n+1} V_i$, and $V_1 = V_2 = V_n = V_{n+1} = k, V_i = k^2$ for all $3 \leq i \leq n-1$. Set $\alpha = (1, 1, 2, \dots, 2, 1, 1) \in \mathbb{N}[I]$, then

$$\mathbb{E}_\alpha = \mathbb{E}_V = \{x \mid x = (x_{13}, x_{23}, x_{43}, \dots, x_{n-1,n-2}, x_{n,n-1}, x_{n+1,n-1}),$$

$$x_{13}, x_{23}, x_{n,n-1}, x_{n+1,n-1} \in M_{21}(k), x_{43}, \dots, x_{n-1,n-2} \in M_{22}(k)\}.$$

Thanks to [DR], we can get $E_1 \oplus E_2 = M(x)$, where $x \in \mathbb{E}_\alpha$ and

$$x_{13} = x_{n,n-1} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, x_{23} = x_{n+1,n-1} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, x_{43} = \dots = x_{n-1,n-2} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

For any $x \in \mathcal{O}_{E_1 \oplus E_2}$, there exists $g = (g_i)_{i \in I} \in GL_\alpha$ such that

$$x_{13} = g_3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_1^{-1}, x_{23} = g_3 \begin{bmatrix} 0 \\ 1 \end{bmatrix} g_2^{-1}, x_{i,i-1} = g_{i-1} g_i^{-1}, 4 \leq i \leq n-1,$$

$$x_{n,n-1} = g_{n-1} \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_n^{-1}, x_{n+1,n-1} = g_{n-1} \begin{bmatrix} 0 \\ 1 \end{bmatrix} g_{n+1}^{-1}.$$

Moreover, we get

$$\begin{bmatrix} x_{13} & x_{23} \end{bmatrix} = x_{43} \cdots x_{n-1,n-2} \cdot \begin{bmatrix} x_{n,n-1} & x_{n+1,n-1} \end{bmatrix} \cdot \begin{bmatrix} g_n g_1^{-1} & 0 \\ 0 & g_{n+1} g_2^{-1} \end{bmatrix},$$

and

$$\begin{bmatrix} x_{n,n-1} & x_{n+1,n-1} \end{bmatrix} = (x_{43} \cdots x_{n-1,n-2})^{-1} \cdot \begin{bmatrix} x_{13} & x_{23} \end{bmatrix} \cdot \begin{bmatrix} g_1 g_n^{-1} & 0 \\ 0 & g_2 g_{n+1}^{-1} \end{bmatrix}.$$

Hence, we have

$$\begin{aligned} \mathcal{O}_{E_1 \oplus E_2} &= \{x \mid x \in \mathbb{E}_\alpha, \begin{bmatrix} x_{13} & x_{23} \end{bmatrix} = x_{43} \cdots x_{n-1,n-2} \cdot \begin{bmatrix} x_{n,n-1} & x_{n+1,n-1} \end{bmatrix} \cdot \begin{bmatrix} \xi_1 & 0 \\ 0 & \xi_2 \end{bmatrix}, \\ &\quad \xi_1 \xi_2 \neq 0, \det(x_{i,i-1}) \neq 0, 4 \leq i \leq n-1\} \\ &= k^* \times k^* \times \{x \mid x = (x_{43}, \dots, x_{n-1,n-2}, x_{n,n-1}, x_{n+1,n-1}), \det(x_{i,i-1}) \neq 0, 4 \leq i \leq n-1 \\ &\quad \det \begin{bmatrix} x_{n,n-1} & x_{n+1,n-1} \end{bmatrix} \neq 0\}. \end{aligned}$$

Let

$$\begin{aligned} X' &= \{x' \mid x' = (x_{43}, \dots, x_{n-1,n-2}, x_{n,n-1}, x_{n+1,n-1}), x_{i,i-1} \in M_{22}(k), 4 \leq i \leq n-1 \\ &\quad x_{n,n-1}, x_{n+1,n-1} \in M_{21}(k)\}. \end{aligned}$$

Then

$$X = \overline{\mathcal{O}}_{E_1 \oplus E_2} = k \times k \times X' = \{x \mid x = (\xi_1, \xi_2, x'), \xi_1, \xi_2 \in k, x' \in X'\} = \mathbb{A}^{4n-10}.$$

The result is clear, now. $\square$

Lemma 4.2.3 *Let $Q = \tilde{E}_6$ as in Fig.4.2.3, let $E_i, i = 1, 2$; $E'_j, j = 1, 2, 3$; $E''_m, m = 1, 2, 3$ be regular simple modules corresponding to three non-homogeneous tubes, respectively. Set $X_1 = \overline{\mathcal{O}}_{E_1 \oplus E_2}$, $X_2 = \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3}$, and $X_3 = \overline{\mathcal{O}}_{E''_1 \oplus E''_2 \oplus E''_3}$. Then $IC_{X_i}(\overline{\mathbb{Q}}_l)$ have the purity property for any $i = 1, 2, 3$.*

Proof. We only prove $i = 1$. The other cases can be proved similarly. Let V be the I -graded vector space $V = \bigoplus_{i=1}^7 V_i$, and $V_1 = V_5 = V_7 = k, V_2 = V_4 = V_6 = k^2, V_3 = k^3$. Set $\alpha = (1, 2, 3, 2, 1, 2, 1) \in \mathbb{N}[I]$, then

$$\begin{aligned} \mathbb{E}_\alpha = \mathbb{E}_V &= \{x \mid x = (x_{12}, x_{23}, x_{43}, x_{54}, x_{63}, x_{76}), \\ &\quad x_{12}, x_{54}, x_{76} \in M_{21}(k), x_{23}, x_{43}, x_{63} \in M_{32}(k)\}. \end{aligned}$$

Thanks to [DR], we can get $E_1 \oplus E_2 = M(x)$, where $x \in \mathbb{E}_\alpha$ and

$$x_{12} = x_{54} = x_{76} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, x_{23} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}, x_{43} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}, x_{63} = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

For any $x \in \mathcal{O}_{E_1 \oplus E_2}$, there exists $g = (g_i)_{i \in I} \in GL_\alpha$ such that

$$\begin{aligned} (4.2.6) \quad x_{12} &= g_2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_1^{-1}, x_{23} = g_3 \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} g_2^{-1}, x_{54} = g_4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_5^{-1}, \\ x_{43} &= g_3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} g_4^{-1}, x_{76} = g_6 \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_7^{-1}, x_{63} = g_3 \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} g_6^{-1}. \end{aligned}$$

Set

$$Z = \{z = (z_0, [\lambda_1 : \lambda_2], z_2, [z_3, z_4, z_5], x) \mid z \in k^2 \times \mathbb{P}^1 \times k^* \times P(k^2 \times k^2 \times k^2) \times \mathbb{E}_\alpha, \\ x_{12}\lambda_1 = z_0\lambda_2, x_{23}z_0 + x_{43}x_{54}z_2 = x_{63}x_{76}, \quad x_{23}z_3 = x_{43}z_4 = x_{63}z_5, \\ \det \begin{bmatrix} z_0 & z_3 \end{bmatrix} \neq 0, \det \begin{bmatrix} x_{12} & z_3 \end{bmatrix} \neq 0, \quad \det \begin{bmatrix} x_{54} & z_4 \end{bmatrix} \neq 0, \quad \det \begin{bmatrix} x_{76} & z_5 \end{bmatrix} \neq 0 \\ \det \begin{bmatrix} x_{23}x_{12} & x_{43}x_{54} & x_{63}z_5 \end{bmatrix} \neq 0, \lambda_1\lambda_2 \neq 0\}.$$

The polynomial map

$$\varphi : Z \longrightarrow \mathcal{O}_{E_1 \oplus E_2}$$

defined by $\varphi(z) = x$. We will show that φ is an isomorphism.

On the one hand, because $z \in Z$, we take

$$g_1 = z_1 = \frac{\lambda_1}{\lambda_2}, g_5 = z_2, g_7 = 1, g_2 = \begin{bmatrix} x_{12}z_1 & z_3 \end{bmatrix}, g_4 = \begin{bmatrix} x_{54}z_2 & z_4 \end{bmatrix}, \\ g_6 = \begin{bmatrix} x_{76} & z_5 \end{bmatrix}, g_3 = \begin{bmatrix} x_{23}x_{12}z_1 & x_{43}x_{54}z_2 & x_{63}z_5 \end{bmatrix}$$

Hence,

$$\begin{aligned} g_2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_1^{-1} &= \begin{bmatrix} x_{12}z_1 & z_3 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} z_1^{-1} = x_{12}, \\ g_3 \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} g_2^{-1} &= \begin{bmatrix} x_{23}x_{12}z_1 & x_{43}x_{54}z_2 & x_{63}z_5 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_{12}z_1 & z_3 \end{bmatrix}^{-1} \\ &= \begin{bmatrix} x_{23}x_{12}z_1 & x_{63}z_5 \end{bmatrix} \begin{bmatrix} x_{12}z_1 & z_3 \end{bmatrix}^{-1} \\ &= \begin{bmatrix} x_{23}x_{12}z_1 & x_{23}z_3 \end{bmatrix} \begin{bmatrix} x_{12}z_1 & z_3 \end{bmatrix}^{-1} \\ &= x_{23} \begin{bmatrix} x_{12}z_1 & z_3 \end{bmatrix} \begin{bmatrix} x_{12}z_1 & z_3 \end{bmatrix}^{-1} \\ &= x_{23}, \\ g_4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_5^{-1} &= g_4 = \begin{bmatrix} x_{54}z_2 & z_4 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} z_2^{-1} = x_{54}, \\ g_3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} g_4^{-1} &= \begin{bmatrix} x_{23}x_{12}z_1 & x_{43}x_{54}z_2 & x_{63}z_5 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_{54}z_2 & z_4 \end{bmatrix}^{-1} \\ &= \begin{bmatrix} x_{43}x_{54}z_2 & x_{43}z_4 \end{bmatrix} \begin{bmatrix} x_{54}z_2 & z_4 \end{bmatrix}^{-1} \\ &= x_{43}, \\ g_6 \begin{bmatrix} 1 \\ 0 \end{bmatrix} &= \begin{bmatrix} x_{76} & z_5 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = x_{76}, \\ g_3 \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} g_6^{-1} &= \begin{bmatrix} x_{23}x_{12}z_1 & x_{43}x_{54}z_2 & x_{63}z_5 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_{76} & z_5 \end{bmatrix}^{-1} \\ &= \begin{bmatrix} x_{63}x_{76} & x_{63}z_5 \end{bmatrix} \begin{bmatrix} x_{76} & z_5 \end{bmatrix}^{-1} \\ &= x_{63}. \end{aligned}$$

So φ is well defined.

On the other hand, we have the inverse map ψ of φ

$$\psi : \mathcal{O}_{E_1 \oplus E_2} \longrightarrow Z, \psi(x) = z,$$

and z is defined in the following.

Since $x \in \mathcal{O}_{E_1 \oplus E_2}$, there exists $(g_i)_{i \in I} \in GL_\alpha$ such that x satisfies (4.2.6). Therefore,

$$(4.2.7) \quad \begin{aligned} x_{23}x_{12}g_1 &= g_3 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad x_{43}x_{54}g_5 = g_3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad x_{63}x_{76}g_7 = g_3 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \\ x_{23}g_2 &= g_3 \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}, \quad x_{43}g_4 = g_3 \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad x_{63}g_6 = g_3 \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}. \end{aligned}$$

Set

$$\frac{\lambda_1}{\lambda_2} = z_1 = \frac{g_1}{g_7}, \quad z_0 = x_{12}z_1, \quad z_2 = \frac{g_5}{g_7}, \quad z_3 = g_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad z_4 = g_4 \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad z_5 = g_6 \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

we have

$$x_{23}x_{12}z_1 + x_{43}x_{54}z_2 = x_{63}x_{76}, \quad x_{23}z_3 = x_{43}z_4 = x_{63}z_5.$$

Based on (4.2.7), we known that $x_{23}x_{12}, x_{43}x_{54}$ is linearly independent over k . Thus z_1 and z_2 is unique determined by x .

Suppose

$$x_{23}z'_3 = x_{43}z'_4 = x_{63}z'_5,$$

and

$$\det \begin{bmatrix} x_{12} & z'_3 \end{bmatrix} \neq 0, \quad \det \begin{bmatrix} x_{54} & z'_4 \end{bmatrix} \neq 0, \quad \det \begin{bmatrix} x_{76} & z'_5 \end{bmatrix} \neq 0, \quad \det \begin{bmatrix} x_{23}x_{12} & x_{43}x_{54} & x_{63}z'_5 \end{bmatrix} \neq 0.$$

By

$$\dim_k C(x_{23}) \cap C(x_{43}) = \dim_k C(x_{43}) \cap C(x_{63}) = \dim_k C(x_{63}) \cap C(x_{23}) = 1$$

we have

$$(4.2.8) \quad x_{23}z'_3 = ax_{23}z_3, \quad x_{43}z'_4 = ax_{43}z_4, \quad x_{63}z'_5 = ax_{63}z_5$$

for some $a \in k^*$.

Since

$$2 = \text{rank}(x_{23}) = \text{rank}(x_{43}) = \text{rank}(x_{63}),$$

(4.2.8) implies

$$z'_3 = az_3, \quad z'_4 = az_4, \quad z'_5 = az_5.$$

Hence, ψ is well defined, and φ is an isomorphism.

Furthermore, let

$$\begin{aligned} X &= \{z = (z_0, [\lambda_1 : \lambda_2], z_2, [z_3, z_4, z_5], x) \mid z \in k^2 \times \mathbb{P}^1 \times k^* \times P(k^2 \times k^2 \times k^2) \times \mathbb{E}_\alpha, \\ &\quad x_{12}\lambda_1 = z_0\lambda_2, x_{23}z_0 + x_{43}x_{54}z_2 = x_{63}x_{76}, \quad x_{23}z_3 = x_{43}z_4 = x_{63}z_5\}. \end{aligned}$$

it is easy to see that $X = \overline{\mathcal{O}}_{E_1 \oplus E_2}$.

We denote by $P(i)$ (resp. $I(i)$) the projective (resp. injective) module corresponding to i for any $i \in I$. Set $Y = \overline{\mathcal{O}}_{\tau^{-2}P(1) \oplus E_2 \oplus I(1)}$. Since

$$\text{Hom}_\Lambda(\tau^{-2}P(1), I(1)) = \text{Hom}_\Lambda(E_2, I(1)) = 0,$$

and

$$\langle \underline{\dim} \tau^{-2}P(1), \underline{\dim} E_2 \rangle = \dim_k \text{Hom}(\tau^{-2}P(1), E_2) = 0,$$

we obtain

$$\dim_k \text{End}(\tau^{-2}P(1) \oplus E_2 \oplus I(1)) = 3, \quad \dim_k X = \dim_k Y + 1.$$

Furthermore, up to isomorphism ψ , we have

$$Y = \{z \mid z \in X, x_{12} = 0, \lambda_2 = 0\}.$$

Let $x' \in \mathbb{E}_\alpha$ such that $x = (x_{12}, 0) + x'$. We define the smooth morphism

$$p : X \longrightarrow Y, \quad p(z) = (z_0, [\lambda_1 : 0], z_2, [z_3, z_4, z_5], x').$$

Since $\tau^{-1}P(1) \oplus E_2 \oplus I(1)$ is aperiodic, $IC_Y(\overline{\mathbb{Q}}_l)$ has the purity property by Theorem 5.4 in [L4]. It is clear that p is smooth with relative dimension 1. Therefore, by Lemma 4.1.4, the statement is proved. $\square$

Lemma 4.2.4 *Let $Q = \tilde{E}_7$ as in Fig.4.2.4, let $E_i, i = 1, 2$; $E'_j, j = 1, 2, 3$; $E''_m, m = 1, 2, 3, 4$ be regular simple objects corresponding to three non-homogeneous tubes, respectively. Set $X_1 = \overline{\mathcal{O}}_{E_1 \oplus E_2}, X_2 = \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3}$, and $X_3 = \overline{\mathcal{O}}_{E''_1 \oplus E''_2 \oplus E''_3 \oplus E''_4}$. Then $IC_{X_i}(\overline{\mathbb{Q}}_l)$ have the purity property for any $i = 1, 2, 3$.*

Proof. We only prove $i = 3$. The other cases can be proved similarly. Let V be the I -graded vector space $V = \bigoplus_{i=1}^8 V_i$, and $V_1 = V_7 = k, V_2 = V_6 = V_8 = k^2, V_3 = V_5 = k^3, V_4 = k^4$. Set $\alpha = (1, 2, 3, 4, 3, 2, 1, 2) \in \mathbb{N}[I]$, then

$$\begin{aligned} \mathbb{E}_\alpha = \mathbb{E}_V &= \{x \mid x = (x_{12}, x_{23}, x_{34}, x_{54}, x_{65}, x_{76}, x_{84}), \\ &\quad x_{12}, x_{76} \in M_{21}(k), x_{23}, x_{65} \in M_{32}(k), x_{34}, x_{54} \in M_{43}(k), x_{84} \in M_{42}(k)\}. \end{aligned}$$

Thanks to [DR], we can get $E''_1 \oplus E''_2 \oplus E''_3 \oplus E''_4 = M(x)$, where $x \in \mathbb{E}_\alpha$ and

$$\begin{aligned} x_{12} = x_{76} &= \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad x_{23} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad x_{65} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad x_{84} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}, \\ x_{34} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \quad x_{54} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}. \end{aligned}$$

For any $x \in \mathcal{O}_{E''_1 \oplus E''_2 \oplus E''_3 \oplus E''_4}$, there exists $(g_i)_{i \in I} \in GL_\alpha$ such that

$$(4.2.8) \quad x_{12} = g_2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_1^{-1}, x_{23} = g_3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} g_2^{-1}, x_{34} = g_4 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} g_3^{-1},$$

$$x_{54} = g_4 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} g_5^{-1}, x_{65} = g_5 \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} g_6^{-1}, x_{76} = g_6 \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_7^{-1},$$

$$x_{84} = g_4 \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} g_8^{-1}.$$

Set

$$Z = \{z = (z_0, [\lambda_1 : \lambda_2], z_2, z_3, z_{41}, z_{42}, z_5, z_6, z_7, x) | z \in k^2 \times \mathbb{P}^1 \times k^2 \times k^3 \times k^2 \times k^2 \times k^3 \times k^2 \times k^* \times \mathbb{E}_\alpha, \\ x_{12}\lambda_1 = z_0\lambda_2, x_{34}x_{23}z_0 = x_{54}z_5, x_{34}x_{23}z_2 = x_{84}z_{41}, x_{34}z_3 = x_{54}x_{65}x_{76}z_7, x_{84}z_{42} = x_{54}x_{65}z_6, \\ \begin{bmatrix} z_2 & : & z_5 \end{bmatrix} \in P(k^2 \times k^3), \begin{bmatrix} z_2 & : & z_{41} \end{bmatrix} \in P(k^2 \times k^2), \begin{bmatrix} z_3 & : & z_7 \end{bmatrix} \in P(k^3 \times k), \\ \begin{bmatrix} z_{42} & : & z_6 \end{bmatrix} \in P(k^2 \times k^2), \begin{bmatrix} z_0 & z_2 \end{bmatrix}, \begin{bmatrix} z_{41} & z_{42} \end{bmatrix}, \begin{bmatrix} x_{12} & z_2 \end{bmatrix}, \begin{bmatrix} x_{76} & z_6 \end{bmatrix} \in GL_2, \\ \begin{bmatrix} x_{23} & z_3 \end{bmatrix}, \begin{bmatrix} z_5 & x_{65} \end{bmatrix} \in GL_3, \lambda_1\lambda_2 \neq 0\}$$

The polynomial map

$$\varphi : Z \longrightarrow \mathcal{O}_{E'_1 \oplus E'_2 \oplus E'_3 \oplus E'_4}$$

defined by $\varphi(z) = x$. We will show that φ is an isomorphism.

On the one hand, we take

$$g_1 = z_1 = \frac{\lambda_1}{\lambda_2}, g_2 = \begin{bmatrix} x_{12}z_1 & z_2 \end{bmatrix}, g_3 = \begin{bmatrix} x_{23}x_{12}z_1 & x_{23}z_2 & z_3 \end{bmatrix}, g_5 = \begin{bmatrix} z_5 & x_{65}x_{76}z_7 & x_{65}z_6 \end{bmatrix}, \\ g_6 = \begin{bmatrix} x_{76}z_7 & z_6 \end{bmatrix}, g_7 = z_7, g_8 = \begin{bmatrix} z_{41} & z_{42} \end{bmatrix}, g_4 = \begin{bmatrix} x_{34}x_{23}x_{12}z_1 & x_{34}x_{23}z_2 & x_{34}z_3 & x_{54}x_{65}z_6 \end{bmatrix}.$$

Thus,

$$g_2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_1^{-1} = \begin{bmatrix} x_{12}z_1 & z_2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} z_1^{-1} = x_{12}, \\ g_3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} g_2^{-1} = \begin{bmatrix} x_{23}x_{12}z_1 & x_{23}z_2 & z_3 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x_{12}z_1 & z_2 \end{bmatrix}^{-1} = x_{23}, \\ g_4 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} g_3^{-1} = \begin{bmatrix} x_{34}x_{23}x_{12}z_1 & x_{34}x_{23}z_2 & x_{34}z_3 & x_{54}x_{65}z_6 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_{23}x_{12}z_1 & x_{23}z_2 & z_3 \end{bmatrix}^{-1} \\ = x_{34}, \\ g_4 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} g_5^{-1} = \begin{bmatrix} x_{34}x_{23}x_{12}z_1 & x_{34}x_{23}z_2 & x_{34}z_3 & x_{54}x_{65}z_6 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} z_5 & x_{65}x_{76}z_7 & x_{65}z_6 \end{bmatrix}^{-1} \\ = \begin{bmatrix} x_{34}x_{23}x_{12}z_1 & x_{34}z_3 & x_{54}x_{65}z_6 \end{bmatrix} \begin{bmatrix} z_5 & x_{65}x_{76}z_7 & x_{65}z_6 \end{bmatrix}^{-1} = x_{54}, \\ g_5 \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} g_6^{-1} = \begin{bmatrix} z_5 & x_{65}x_{76}z_7 & x_{65}z_6 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_{76}z_7 & z_6 \end{bmatrix}^{-1} = x_{65}, \\ g_6 \begin{bmatrix} 1 \\ 0 \end{bmatrix} g_7^{-1} = \begin{bmatrix} x_{76}z_7 & z_6 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = x_{76}, \\ g_4 \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} g_8^{-1} = \begin{bmatrix} x_{34}x_{23}x_{12}z_1 & x_{34}x_{23}z_2 & x_{34}z_3 & x_{54}x_{65}z_6 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} z_{41} & z_{42} \end{bmatrix}^{-1} = x_{84},$$

that is, φ is well defined.

On the other hand, we get the inverse map ψ of φ

$$\psi : \mathcal{O}_{E'_1 \oplus E'_2 \oplus E'_3 \oplus E'_4} \longrightarrow Z, \psi(x) = z,$$

and z is defined in the following.

Since $x \in \mathcal{O}_{E'_1 \oplus E'_2 \oplus E'_3 \oplus E'_4}$, there exists $(g_i)_{i \in I} \in GL_\alpha$ such that x satisfies (4.2.8). Therefore,

$$(4.2.9) \quad \begin{aligned} x_{34}x_{23}x_{12}g_1 &= x_{54}g_5 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = g_4 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad x_{34}x_{23}g_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} = x_{84}g_8 \begin{bmatrix} 1 \\ 0 \end{bmatrix} = g_4 \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \\ x_{34}g_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} &= x_{54}x_{65}x_{76}g_7 = g_4 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \quad x_{84}g_8 \begin{bmatrix} 0 \\ 1 \end{bmatrix} = x_{54}x_{65}g_6 \begin{bmatrix} 0 \\ 1 \end{bmatrix} = g_4 \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}. \end{aligned}$$

Set

$$\begin{aligned} \frac{\lambda_1}{\lambda_2} &= z_1 = g_1, \quad z_0 = x_{12}z_1, \quad z_2 = g_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad z_3 = g_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \quad z_{41} = g_8 \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad z_{42} = g_8 \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \\ z_5 &= g_5 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad z_6 = g_6 \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad z_7 = g_7. \end{aligned}$$

It is easy to see that $z = (z_0, [\lambda_1 : \lambda_2], z_2, z_3, z_{41}, z_{42}, z_5, z_6, z_7, x) \in Z$.

Suppose $z' = (z'_0, [\lambda'_1 : \lambda'_2], z'_2, z'_3, z'_{41}, z'_{42}, z'_5, z'_6, z'_7, x) \in Z$, we have

$$(4.2.10) \quad \begin{aligned} x_{34}x_{23}z_0 &= x_{54}z_5, \quad x_{34}x_{23}z_2 = x_{84}z_{41}, \quad x_{34}z_3 = x_{54}x_{65}x_{76}z_7, \quad x_{84}z_{42} = x_{54}x_{65}z_6, \\ x_{34}x_{23}z'_0 &= x_{54}z'_5, \quad x_{34}x_{23}z'_2 = x_{84}z'_{41}, \quad x_{34}z'_3 = x_{54}x_{65}x_{76}z'_7, \quad x_{84}z'_{42} = x_{54}x_{65}z'_6. \end{aligned}$$

Since $\dim_k C(x_{54}) \cap C(x_{34}x_{23}) = 1$, we have $x_{54}z'_5 = \lambda x_{54}z_5$ for some λ . Because $\text{rank}(x_{54}) = 3$, there exists a matrix y_{54} such that $y_{54}x_{54} = I_3$. Thus, $z_5 = \lambda z'_5$, that is, $[z_0 : z_5] \in P(k^2 \times k^3)$.

Based on (4.2.8), we have

$$C(x_{34}x_{23}) = C(g_4 \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}) \text{ and } C(x_{84}) = C(g_4 \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}).$$

Since

$$C(x_{84}) \cap C(x_{34}x_{23}) = C(g_4 \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}),$$

we obtain $z_2 = \mu z'_2$ and $z_{41} = \mu z'_{41}$ for some $\mu \in k^*$, that is $[z_2 : z_{41}] \in P(k^2 \times k^2)$. In the same way, we get $[z_3 : z_7] \in P(k^3 \times k)$, $[z_{42} : z_6] \in P(k^2 \times k^2)$.

Hence, ψ is well defined, and φ is an isomorphism.

Let

$$\begin{aligned} X = \{z = (z_0, [\lambda_1 : \lambda_2], z_2, z_3, z_{41}, z_{42}, z_5, z_6, z_7, x) \mid & z \in k^2 \times \mathbb{P}^1 \times k^2 \times k^3 \times k^2 \times k^2 \times k^3 \times k^2 \times k \times \mathbb{E}_\alpha, \\ & x_{12}\lambda_1 = z_0\lambda_2, \quad x_{34}x_{23}z_0 = x_{54}z_5, \quad x_{34}x_{23}z_2 = x_{84}z_{41}, \quad x_{34}z_3 = x_{54}x_{65}x_{76}z_7, \quad x_{84}z_{42} = x_{54}x_{65}z_6, \\ & \begin{bmatrix} z_0 & : & z_5 \end{bmatrix} \in P(k^2 \times k^3), \quad \begin{bmatrix} z_2 & : & z_{41} \end{bmatrix} \in P(k^2 \times k^2), \quad \begin{bmatrix} z_3 & : & z_7 \end{bmatrix} \in P(k^3 \times k), \\ & \begin{bmatrix} z_{42} & : & z_6 \end{bmatrix} \in P(k^2 \times k^2), \} \end{aligned}$$

Then $X = \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3 \oplus E'_4}$.

Set $Y = \overline{\mathcal{O}}_{\tau^{-2}P(7) \oplus E_2'' \oplus E_3'' \oplus E_4'' \oplus I(1)}$. Since

$$\text{Hom}_{\Lambda}(\tau^{-2}P(7), I(1)) = \text{Hom}_{\Lambda}(E_i'', I(1)) = 0, i = 2, 3, 4$$

$$\langle \underline{\dim} \tau^{-2}P(7), \underline{\dim} E_i'' \rangle = \dim_k \text{Hom}_{\Lambda}(\tau^{-2}P(7), E_i'') = 0, i = 2, 3, 4,$$

we have

$$\dim_k \text{End}_{\Lambda}(\tau^{-2}P(7) \oplus E_2'' \oplus E_3'' \oplus E_4'' \oplus I(1)) = 5, \dim X = \dim Y + 1.$$

Furthermore, up to isomorphism φ , we have

$$Y = \{z | z \in X, x_{12} = 0, \lambda_2 = 0\}$$

Let $x' \in \mathbb{E}_{\alpha}$ such that $x = (x_{12}, 0) + x'$, and define the smooth morphism

$$p : X \longrightarrow Y, p(z) = (z_0, [\lambda_1 : 0], z_2, z_3, z_4, z_5, z_6, z_7, x').$$

Thus, Lemma 4.2.4 follows from the lemma 4.1.4. $\square$

Lemma 4.2.5 *Let $Q = \widetilde{E}_8$ as in Fig.4.2.5, let $E_i, i = 1, 2; E'_j, j = 1, 2, 3; E''_m, m = 1, 2, 3, 4, 5$ be regular simple objects corresponding to three non-homogeneous tubes, respectively. Set $X_1 = \overline{\mathcal{O}}_{E_1 \oplus E_2}, X_2 = \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3}$, and $X_3 = \overline{\mathcal{O}}_{E''_1 \oplus E''_2 \oplus E''_3 \oplus E''_4 \oplus E''_5}$. Then $IC_{X_i}(\overline{\mathbb{Q}}_l)$ have the purity property for any $i = 1, 2, 3$.*

Proof. We only prove $i = 1$. The other cases can be proved similarly. Let V be the I -graded vector space $V = \bigoplus_{i=1}^9 V_i$, and $V_1 = V_7 = k^2, V_2 = V_5 = k^4, V_3 = k^6, V_4 = k^5, V_6 = V_9 = k^3, V_8 = k$. Set $\alpha = (2, 4, 6, 5, 4, 3, 2, 1, 3) \in \mathbb{N}[I]$, then

$$\begin{aligned} \mathbb{E}_{\alpha} = \mathbb{E}_V &= \{x | x = (x_{12}, x_{23}, x_{43}, x_{54}, x_{65}, x_{76}, x_{87}, x_{93}), \\ &\quad x_{12} \in M_{42}(k), x_{23} \in M_{64}(k), x_{43} \in M_{65}(k), x_{54} \in M_{54}(k), x_{65} \in M_{43}(k), x_{76} \in M_{32}(k), \\ &\quad x_{87} \in M_{21}(k), x_{93} \in M_{63}(k)\}. \end{aligned}$$

Thanks to [DR], we can get $E_1 \oplus E_2 = M(x)$, where $x \in \mathbb{E}_{\alpha}$ and

$$\begin{aligned} x_{12} &= \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}, \quad x_{23} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad x_{43} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad x_{54} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \\ x_{65} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad x_{76} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad x_{87} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad x_{93} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1. \end{bmatrix}. \end{aligned}$$

For any $x \in \mathcal{O}_{E_1 \oplus E_2}$, there exists $(g_i)_{i \in I} \in GL_\alpha$ such that

$$(4.2.11) \quad \begin{aligned} x_{12} &= g_2 \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} g_1^{-1}, x_{23} = g_3 \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} g_2^{-1}, x_{43} = g_3 \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} g_4^{-1}, \\ x_{54} &= g_4 \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} g_5^{-1}, x_{65} = g_5 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} g_6^{-1}, x_{76} = g_6 \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix} g_7^{-1}, \\ x_{87} &= g_7 \begin{bmatrix} 0 \\ 1 \end{bmatrix} g_8^{-1}, x_{93} = g_3 \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix} g_9^{-1}. \end{aligned}$$

Set

$$\begin{aligned} Z = \{ & z = (z_{11}, z_{12}, z_2, z_3, z_4, z_5, z_6, z_7, z_8, [\lambda_1 : \lambda_2], z_{91}, z_{92}, z_{93}, x) | \\ & z \in k^2 \times k^2 \times k^4 \times k^4 \times k^5 \times k^4 \times k^3 \times k^2 \times k^2 \times \mathbb{P}^1 \times k^3 \times k^3 \times k^3 \times \mathbb{E}_\alpha, \ x_{87}\lambda_1 = z_8\lambda_2, \\ & x_{23}z_2 = x_{43}x_{54}z_5, \ x_{23}z_3 = x_{43}x_{54}x_{65}z_6, \ x_{93}z_{91} = x_{43}x_{54}x_{65}x_{76}z_7 + x_{23}z_2, \\ & x_{93}z_{92} = x_{23}x_{12}z_{11} + x_{43}x_{54}z_5, \ x_{23}x_{12}z_{12} = x_{43}z_4, \\ & x_{93}z_{93} = x_{43}x_{54}x_{65}x_{76}z_8 + x_{23}z_3 + x_{23}x_{12}z_{12}, \\ & [z_3 : z_6] \in P(k^4 \times k^3), [z_{12} : z_4] \in P(k^2 \times k^5), \begin{bmatrix} z_{11} & z_{12} \end{bmatrix}, \begin{bmatrix} z_7 & z_8 \end{bmatrix} \in GL_2, \\ & \begin{bmatrix} x_{76}z_7 & x_{76}z_8 & z_6 \end{bmatrix} \in GL_3 \begin{bmatrix} z_2 & x_{12}z_{11} & z_3 & x_{12}z_{12} \end{bmatrix}, \\ & \begin{bmatrix} x_{65}x_{76}z_7 & z_5 & x_{65}x_{76}z_8 & x_{65}z_6 \end{bmatrix} \in GL_4 \\ & \begin{bmatrix} z_{91} & z_{92} & z_{93} \end{bmatrix} \in GL_3, \begin{bmatrix} x_{54}x_{65}x_{76}z_7 & x_{54}z_5 & x_{54}x_{65}x_{76}z_8 & x_{54}x_{65}z_6 & z_4 \end{bmatrix} \in GL_5, \ \lambda_1\lambda_2 \neq 0 \} \end{aligned}$$

The polynomial map

$$\varphi : Z \longrightarrow \mathcal{O}_{E_1 \oplus E_2}$$

defined by $\varphi(z) = x$.

If we take

$$\begin{aligned} g_1 &= \begin{bmatrix} z_{11} & z_{12} \end{bmatrix}, \ g_2 = \begin{bmatrix} z_2 & x_{12}z_{11} & z_3 & x_{12}z_{12} \end{bmatrix}, \\ g_3 &= \begin{bmatrix} x_{43}x_{54}x_{65}x_{76}z_7 & x_{43}x_{54}z_5 & x_{23}x_{12}z_{11} & x_{43}x_{54}x_{65}x_{76}z_8 & x_{43}x_{54}x_{65}z_6 & x_{43}z_4 \end{bmatrix}, \\ g_4 &= \begin{bmatrix} x_{54}x_{65}x_{76}z_7 & x_{54}z_5 & x_{54}x_{65}x_{76}z_8 & x_{54}x_{65}z_6 & z_4 \end{bmatrix}, \ g_5 = \begin{bmatrix} x_{65}x_{76}z_7 & z_5 & x_{65}x_{76}z_8 & x_{65}z_6 \end{bmatrix}, \\ g_6 &= \begin{bmatrix} x_{76}z_7 & x_{76}z_8 & z_6 \end{bmatrix}, \ g_7 = \begin{bmatrix} z_7 & z_8 \end{bmatrix}, \end{aligned}$$

one may get $x \in \mathbb{E}_\alpha$ satisfies (4.2.11), that is, φ is well defined.

On the other hand, we have the inverse map ψ of φ

$$\psi : \mathcal{O}_{E_1 \oplus E_2} \longrightarrow Z, \psi(x) = z,$$

and z is defined as in the following.

Since $x \in \mathcal{O}_{E_1 \oplus E_2}$, there exists $(g_i)_i \in GL_{\dim E_1 \oplus E_2}$ such that x satisfies (4.2.11). Moreover,

$$\begin{aligned}
 (4.2.12) \quad & x_{23}g_2 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} = x_{43}x_{54}g_5 \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad x_{23}g_2 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} = x_{43}x_{54}x_{65}g_6 \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \\
 & x_{23}x_{12}g_1 \begin{bmatrix} 0 \\ 1 \end{bmatrix} = x_{43}g_4 \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \quad x_{93}g_9 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = x_{43}x_{54}x_{65}x_{76}g_7 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_{23}g_2 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \\
 & x_{93}g_9 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = x_{23}x_{12}g_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_{43}x_{54}g_5 \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \\
 & x_{93}g_9 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = x_{43}x_{54}x_{65}x_{76}x_{87}g_8 + x_{23}g_2 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + x_{23}g_2 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + x_{23}x_{12}g_1 \begin{bmatrix} 0 \\ 1 \end{bmatrix}.
 \end{aligned}$$

Set

$$\begin{aligned}
 z_{11} &= g_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad z_{12} = g_1 \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad z_2 = g_2 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad z_3 = g_2 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \quad z_4 = g_4 \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \quad z_5 = g_5 \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \\
 z_6 &= g_6 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \quad z_7 = g_7 \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad z_8 = x_{87}g_8, \quad \frac{\lambda_1}{\lambda_2} = g_8,
 \end{aligned}$$

then we have $z = (z_{11}, z_{12}, z_2, z_3, z_4, z_5, z_6, z_7, z_8, [\lambda_1 : \lambda_2], z_{91}, z_{92}, z_{93}, x) \in Z$.

Suppose $z' = (z'_{11}, z'_{12}, z'_2, z'_3, z'_4, z'_5, z'_6, z'_7, z'_8, [\lambda'_1 : \lambda'_2], z'_{91}, z'_{92}, z'_{93}, x) \in Z$, then we also have

$$\begin{aligned}
 (4.2.13) \quad & x_{23}z'_2 = x_{43}x_{54}z'_5, \quad x_{23}z'_3 = x_{43}x_{54}x_{65}z'_6, \quad x_{87}\lambda'_1 = z'_8\lambda'_2, \\
 & x_{93}z'_{91} = x_{43}x_{54}x_{65}x_{76}z'_7 + x_{23}z'_2, \quad x_{93}z'_{92} = x_{23}x_{12}z'_{11} + x_{43}x_{54}z'_5, \\
 & x_{23}x_{12}z'_{12} = x_{43}z'_4, \quad x_{93}z'_{93} = x_{43}x_{54}x_{65}x_{76}z'_8 + x_{23}z'_3 + x_{23}x_{12}z'_{12}.
 \end{aligned}$$

Based on (4.2.11), we obtain

$$C(x_{23}) = C(g_3 \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}), \quad \text{and} \quad C(x_{43}x_{54}x_{65}) = C(g_3 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}),$$

$$C(x_{23}x_{12}) = C(g_3 \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}) \text{ and } C(x_{43}) = C(g_3 \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix})$$

Therefore,

$$C(x_{43}x_{54}x_{65}) \cap C(x_{23}) = C(x_{23}x_{12}) \cap C(x_{43}) = C(g_3 \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}).$$

It implies that $z'_3 = \mu z_3$, $z'_6 = \mu z_6$, $z'_{12} = \nu z_{12}$, and $z'_4 = \nu z_4$ for some μ and ν , that is, $[z_3 : z_6] \in P(k^4 \times k^3)$ and $[z_{12} : z_4] \in P(k^2 \times k^5)$.

Hence ψ is well defined, and φ is an isomorphism.

Let

$$\begin{aligned} X = \{z = (z_{11}, z_{12}, z_2, z_3, z_4, z_5, z_6, z_7, z_8, [\lambda_1 : \lambda_2], z_{91}, z_{92}, z_{93}, x) \mid \\ z \in k^2 \times k^2 \times k^4 \times k^4 \times k^5 \times k^4 \times k^3 \times k^2 \times k^2 \times \mathbb{P}^1 \times k^3 \times k^3 \times k^3 \times \mathbb{E}_\alpha, \quad x_{23}z_2 = x_{43}x_{54}z_5, \\ x_{23}z_3 = x_{43}x_{54}x_{65}z_6, \quad x_{93}z_{91} = x_{43}x_{54}x_{65}x_{76}z_7 + x_{23}z_2, \quad x_{93}z_{92} = x_{23}x_{12}z_{11} + x_{43}x_{54}z_5, \\ x_{23}x_{12}z_{12} = x_{43}z_4, \quad x_{93}z_{93} = x_{43}x_{54}x_{65}x_{76}x_{87}z_8 + x_{23}z_3 + x_{23}x_{12}z_{12}, \quad x_{87}\lambda_1 = z_8\lambda_2, \\ [z_3 : z_6] \in P(k^4 \times k^3), [z_{12} : z_4] \in P(k^2 \times k^5)\} \end{aligned}$$

then we get $X = \overline{\mathcal{O}}_{E_1 \oplus E_2}$.

Let P be projective with dimension vector (123322101), and set $Y = \overline{\mathcal{O}}_{P \oplus E_1 \oplus I(8)}$.

Since

$$\begin{aligned} \text{Hom}_\Lambda(P, I(8)) &= \text{Hom}_\Lambda(E_1, I(8)) = 0, \\ \langle \underline{\dim} P, \underline{\dim} E_1 \rangle &= \text{Hom}_\Lambda(P, E_1) = 0, \end{aligned}$$

we get

$$\dim \text{End}_\Lambda(P \oplus E_1 \oplus I(8)) = 3, \dim X = \dim Y + 1.$$

Furthermore, up to isomorphism φ , we have

$$Y = \{z \mid z \in X, x_{87} = \lambda_2 = 0\}.$$

Let $x' \in \mathbb{E}_\alpha$ such that $x = (0, x_{87}, 0) + x'$ and define

$$p : X \longrightarrow Y, p(z) = (z_{11}, z_{12}, z_2, z_3, z_4, z_5, z_6, z_7, z_8, [\lambda_1 : 0], z_{91}, z_{92}, z_{93}, x').$$

Applying Lemma 4.1.4, the proof is complete. $\square$

Furthermore, in the same way, we can prove that the closure of orbits of semi-simple objects have the purity property.

In order to prove Theorem 4.1.5, we not only need to discuss the closure of semi-simple objects in $\mathcal{T}_i$ which have purity property, but also need to study the fibres of p_3 .

5. The Fibres of p_3

Let $\mathcal{P}_\gamma$ be a set of Λ modules of dimension vector γ , up to isomorphism. From the definition of p_3 in 1.3, it follows that $p_3^{-1}(L) = \cup_{N \in \mathcal{P}_\beta, N \subseteq L} \mathbf{Z}_{L, M, N}$. Thus we need to discuss some properties of the variety $\mathbf{Z}_{L, M, N}$. In fact, we know that g_{MN}^L is the number of rational points of varieties

$\mathbf{Z}_{L,M,N}$. Set $Z_L = \{(M, N) | (M, N) \in \mathcal{P}_\alpha \times \mathcal{P}_\beta, N \subseteq L, \text{ and } L/N \cong M\}$. If Z_L has only finitely many elements, then $p^{-1}(L)$ has a natural stratification with strata $\mathbf{Z}_{L,M,N}$ indexed by $(M, N) \in Z_L$.

For the convenience of discussions below, we need to introduce some notations about the BGP reflection functors (see[BGP] or[DR]).

We define $\sigma_i Q$ to be the quiver obtained from Q by reversing the direction of every arrow connected to the vertex i . If i is a sink of Q , σ_i^+ is defined as follows:

$$\sigma_i^+ : \text{mod } \Lambda \longrightarrow \text{mod } \sigma_i \Lambda,$$

where $\Lambda = \mathbb{F}_q(Q)$ (resp. $\Lambda = k(Q)$) and $\sigma_i \Lambda = \mathbb{F}_q(\sigma_i Q)$ (resp. $\sigma_i \Lambda = k(\sigma_i Q)$) are path algebras. Therefore σ_i^+ is an exact functor on the full subcategory $\text{mod } \Lambda(i)$ of $\text{mod } \Lambda$ which do not have S_i as a direct summand, and induce quasi-inverse equivalence between $\text{mod } \Lambda(i)$ and the full subcategory $\text{mod } \sigma_i \Lambda(i)$ which do not have direct summand isomorphic to the simple injective module S_i .

According to the notation of Hall polynomial (see [R3]), we use $|\cdot|$ to denote the ordinary cardinality of a finite set.

Lemma 5.1 *Let M, M_1, M_2 and N be Λ -modules and let N (resp. M_2) be a pre-projective (resp. regular). If $\text{Hom}_\Lambda(M_2, M_1) = 0$, then*

$$g_{M_1 \oplus M_2, N}^{M \oplus M_2} = g_{M_1, N}^M \frac{|\text{Hom}_\Lambda(M, M_2)|}{|\text{Hom}_\Lambda(M_1, M_2)|}.$$

Proof. Let X, Y and L be Λ -modules, and set

$$W(X, Y; L) = \{(f, g) \in \text{Hom}_\Lambda(X, L) \times \text{Hom}_\Lambda(L, Y) | \\ 0 \longrightarrow X \longrightarrow L \longrightarrow Y \longrightarrow 0 \text{ is a short exact sequence}\}.$$

The action of $\text{Aut}(X) \times \text{Aut}(Y)$ on $W(X, Y; L)$ is defined by

$$(a, c)(f, g) = (fa, c^{-1}g).$$

It induces the orbit space

$$V(X, Y; L) = \{(f, g)^\wedge | (f, g) \in W(X, Y; L)\}.$$

Then $|V(X, Y; L)| = g_{YX}^L$.

Note that the actions of $\text{Aut}(N) \times \text{Aut}(M_1 \oplus M_2)$ and $\text{Aut}(N) \times \text{Aut}(M_1)$ on $W(N, M_1 \oplus M_2; M \oplus M_2)$ and $W(N, M_1; M)$, respectively, are free. So we have

$$(5.1) \quad |V(N, M_1; M)| = \frac{|W(N, M_1; M)|}{|\text{Aut}(N)||\text{Aut}(M_1)|}, \text{ and} \\ |V(N, M_1 \oplus M_2; M \oplus M_2)| = \frac{|W(N, M_1 \oplus M_2; M \oplus M_2)|}{|\text{Aut}(N)||\text{Aut}(M_1 \oplus M_2)|}.$$

If $(\begin{bmatrix} f_1 \\ f_2 \end{bmatrix}, \begin{bmatrix} g_{11} & 0 \\ g_{21} & g_{22} \end{bmatrix}) \in W(N, M_1 \oplus M_2; M \oplus M_2)$, we have g_{22} to be invertible and $f_2 = -g_{22}^{-1}g_{21}f_1$.

Consider the map

$$\varphi : W(N, M_1 \oplus M_2; M \oplus M_2) \longrightarrow W(N, M_1; M) \times \text{Hom}_\Lambda(M, M_2) \times \text{Aut}(M_2), \\ (\begin{bmatrix} f_1 \\ -g_{22}^{-1}g_{21}f_1 \end{bmatrix}, \begin{bmatrix} g_{11} & 0 \\ g_{21} & g_{22} \end{bmatrix}) \longmapsto (f_1, g_{11}, g_{21}, g_{22}).$$

It is easy to see that the inverse map ψ of φ is

$$\begin{aligned} \varphi : W(N, M_1; M) \times \text{Hom}_\Lambda(M, M_2) \times \text{Aut}(M_2) &\longrightarrow W(N, M_1 \oplus M_2; M \oplus M_2), \\ (f_1, g_{11}, g_{21}, g_{22}) &\longmapsto \left(\begin{bmatrix} f_1 \\ -g_{22}^{-1}g_{21}f_1 \end{bmatrix}, \begin{bmatrix} g_{11} & 0 \\ g_{21} & g_{22} \end{bmatrix} \right). \end{aligned}$$

Thus, φ is an isomorphism. Therefore, we have

$$(5.2) \quad |W(N, M_1 \oplus M_2; M \oplus M_2)| = |W(N, M_1; M)| |\text{Hom}_\Lambda(M, M_2) \times \text{Aut}(M_2)|$$

Hence the proof follows from (5.1) and (5.2). $\square$

Proposition 5.2 *Let S be a simple projective modules of kQ (except $Q = \tilde{E}_8$) corresponding to an unique sink point, P a preprojective, and let M be a regular semi-simple objects in $\mathcal{T}_i$ for some $i, 1 \leq i \leq l$. Then $(p_3|_{\overline{\mathcal{O}}_M \times \overline{\mathcal{O}}_S})^{-1}(P \oplus L)$ have purity property.*

Proof. If $g_{MS}^{P \oplus L} \neq 0$, we have

$$0 \longrightarrow S \xrightarrow{f} P \oplus L \xrightarrow{\begin{bmatrix} g_1 \\ g_2 \end{bmatrix}^t} M \longrightarrow 0.$$

According to the representation theory of quivers, we know that $L \in \mathcal{T}_i$. First, we claim that L is a regular semi-simple submodule of M . Obviously, g_2 is a monomorphism. Let, otherwise, $\text{rad}_{\mathcal{T}_i}(L)$ be the regular radical of L in full sub-category $\mathcal{T}_i$ of $\Lambda - \text{mod}$, then we have $g_2(\text{rad}_{\mathcal{T}_i}(L)) = 0$ and $\text{rad}_{\mathcal{T}_i}(L) \subseteq \text{Im}(f)$. But $\text{Hom}_\Lambda(\text{rad}_{\mathcal{T}_i}(L), S) = 0$, a contradiction.

Because of M and L to be semi-simple in $\mathcal{T}_i$. Let l_2 be a morphism from M to L such that $l_2 g_2 = \text{id}_L$. Then $M = M_1 \oplus \text{Im}(g_2)$, where $M_1 = \{x - g_2 l_2(x) | x \in M\}$.

Since

$$M \cong M_1 \oplus \text{Im}(g_2),$$

g_{22} is an isomorphism and there are short exact sequences

$$0 \longrightarrow S \xrightarrow{\begin{bmatrix} f_1 \\ -g_{22}^{-1}g_{21}f_1 \end{bmatrix}} P \oplus L \xrightarrow{\begin{bmatrix} g_{11} & 0 \\ g_{21} & g_{22} \end{bmatrix}} M_1 \oplus \text{Im}(g_2) \longrightarrow 0$$

and

$$0 \longrightarrow S \xrightarrow{f_1} P \xrightarrow{g_{11}} M_1 \longrightarrow 0.$$

For any

$$\left(\begin{bmatrix} f_1 \\ f_2 \end{bmatrix}, \begin{bmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{bmatrix} \right) \in W(S, M_1 \oplus L; P \oplus L).$$

Because of $\begin{bmatrix} g_{12} \\ g_{22} \end{bmatrix}$ to be a split monomorphism, there is a morphism $\begin{bmatrix} l_{21} & l_{22} \end{bmatrix}$ from $M_1 \oplus L$ to L such that

$$\begin{bmatrix} l_{21} & l_{22} \end{bmatrix} \begin{bmatrix} g_{12} \\ g_{22} \end{bmatrix} = \text{id}_L.$$

Without loss of generality, we may assume that l_{22} is an invertible. Otherwise, g_{12} is a split monomorphism, it implies $M_1 = M_2 \oplus L$. We then can get the same argument.

Since

$$\begin{bmatrix} \text{id}_{M_1} & -g_{12} \\ 0 & \text{id}_L \end{bmatrix} \begin{bmatrix} \text{id}_{M_1} & 0 \\ l_{21} & l_{22} \end{bmatrix} \begin{bmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{bmatrix} = \begin{bmatrix} g_{11} - g_{12}(l_{21}g_{11} + l_{22}g_{21}) & 0 \\ l_{21}g_{11} + l_{22}g_{21} & \text{id}_L \end{bmatrix},$$

we have $c = \begin{bmatrix} id_{M_1} & -g_{12} \\ 0 & id_L \end{bmatrix} \begin{bmatrix} id_{M_1} & 0 \\ l_{21} & l_{22} \end{bmatrix} \in Aut_\Lambda(M_1 \oplus L)$ such that

$$c \begin{bmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{bmatrix} = \begin{bmatrix} g'_{11} & 0 \\ g'_{21} & 1 \end{bmatrix}.$$

Thus

$$V(S, M_1 \oplus L; P \oplus L) = \{ \left(\begin{bmatrix} f_1 \\ -g_{21}f_1 \end{bmatrix}, \begin{bmatrix} g_{11} & 0 \\ g_{21} & 1 \end{bmatrix} \right)^\wedge | \left(\begin{bmatrix} f_1 \\ -g_{21}f_1 \end{bmatrix}, \begin{bmatrix} g_{11} & 0 \\ g_{21} & 1 \end{bmatrix} \right) \in W(S, M_1 \oplus L; P \oplus L) \}.$$

Consider the map

$$\varphi : V(S, M_1 \oplus L; P \oplus L) \longrightarrow V(S, M_1; P)$$

sending $\left(\begin{bmatrix} f_1 \\ -g_{21}f_1 \end{bmatrix}, \begin{bmatrix} g_{11} & 0 \\ g_{21} & 1 \end{bmatrix} \right)^\wedge$ to $(f_1, g_{11})^\wedge$. Let $V(g_{11}) = \{hg_{11} | h \in Hom_\Lambda(M_1, L)\}$. Then $V(g_{11})$ is a k -vector subspace of $Hom_\Lambda(P, L)$. Because of g_{11} to be epimorphism, we have $V(g_{11}) \cong Hom_\Lambda(M_1, L)$ as k -space. Thereby

$$(A) \left\{ \begin{array}{l} \varphi^{-1}((f_1, g_{11})^\wedge) = Hom_\Lambda(P, L)/V(g_{11}) = \mathbb{A}^{dim_k Hom_\Lambda(P, L)/V(g_{11})}, \\ |V(S, M_1 \oplus L; P \oplus L)| = |V(S, M_1; P)| \times |\mathbb{A}^{dim_k Hom_\Lambda(P, L)/V(g_{11})}|. \end{array} \right.$$

The rest of the proof can be divided into several cases:

Case 1 : $Q = \tilde{A}_n$.

Let $E_i, 1 \leq i \leq n$ be simple objects in the full non-homogeneous subcategory of $\Lambda - \text{mod}$ corresponding to the dimension vectors listed in Fig 4.2.1. Since there is only simple object E_n such that $Ext^1(E_n, S) \neq 0$, by Lemma 5.1, we only need to discuss the case $N = \oplus mE_n$.

Because $P(n) \oplus \oplus (m-1)E_n$ is an unique non-trivial extension of S by N , it is easy to see that $g_{mE_n, S}^{P(n) \oplus \oplus (m-1)E_n} = 1$.

Set $M = \oplus a_n E_n \oplus \oplus_{i \neq n} a_i E_i$. Since

$$Hom_\Lambda(\oplus_{i \neq n} a_i E_i, \oplus a_n E_n) = Hom_\Lambda(\oplus a_n E_n, \oplus_{i \neq n} a_i E_i) = 0,$$

we obtain, by Lemma 5.1,

$$\begin{aligned} g_{\oplus a_n E_n \oplus \oplus_{i \neq n} a_i E_i, S}^{P(n) \oplus (a_n - 1)E_n \oplus (\oplus_{i \neq n} a_i E_i)} &= g_{\oplus a_n E_n, S}^{P(n) \oplus (a_n - 1)E_n} |Hom_\Lambda(P(n) \oplus (a_n - 1)E_n, \oplus_{i \neq n} a_i E_i)| \\ &= |Hom_\Lambda(P(n), \oplus_{i \neq n} a_i E_i)| \end{aligned}$$

Moreover,

$$\begin{aligned} \mathbf{Z}_{P(n) \oplus (a_n - 1)E_n \oplus (\oplus_{i \neq n} a_i E_i), M, S} &= Hom_\Lambda(P(n), \oplus_{i \neq n} a_i E_i), \\ (p_3|_{\overline{\mathcal{O}}_M \times \overline{\mathcal{O}}_S})^{-1}(P(n) \oplus (a_n - 1)E_n \oplus (\oplus_{i \neq n} a_i E_i)) &= Hom_\Lambda(P(n), \oplus_{i \neq n} a_i E_i). \end{aligned}$$

Therefore $\mathbf{Z}_{P(n) \oplus (a_n - 1)E_n \oplus (\oplus_{i \neq n} a_i E_i), M, S}$ has purity property.

Case 2 : $Q = \tilde{D}_n, n \geq 4$.

We only consider $\mathcal{T}_1$ in Fig 4.2.2. The other cases can be proved similarly. Let $E_i, i = 1, 2$ be simple objects in the full subcategory $\mathcal{T}_1$ of $\Lambda - \text{mod}$ corresponding to the dimension vectors listed in Fig 4.2.2.

Since

$$dim_k Ext^1(E_i, S) = -\langle \underline{\dim} E_i, \underline{\dim} S \rangle = 1,$$

we have

$$dim_k Ext^1(\oplus mE_i, S) = m.$$

Moreover,

$$\xi : 0 \longrightarrow S \xrightarrow{f_\xi = \begin{bmatrix} 1 \\ -1 \end{bmatrix}} P(1) \oplus P(n) \xrightarrow{g_\xi = \begin{bmatrix} g_1 & g_2 \end{bmatrix}} E_1 \longrightarrow 0$$

and

$$\eta : 0 \longrightarrow S \xrightarrow{\begin{bmatrix} 1 \\ -1 \end{bmatrix}} P(2) \oplus P(n+1) \longrightarrow E_2 \longrightarrow 0$$

are unique non-trivial extensions of S by E_1 and E_2 , respectively.

Based on the claim above, let M be a non-trivial extension of S by $\oplus mE_1 \oplus \oplus rE_2$, it is of the form $P \oplus (\oplus(m-a)E_1 \oplus \oplus(r-b)E_2)$ for some $0 \leq a \leq m, 0 \leq b \leq r$, where P is a pre-projective.

Obviously, the non-trivial extension of S by $\oplus mE_1$ (resp. $\oplus rE_2$) has unique form $P(1) \oplus P(n) \oplus \oplus(m-1)E_1$ (resp. $P(2) \oplus P(n+1) \oplus \oplus(r-1)E_2$).

Since

$$|Ext_\Lambda^1(\oplus mE_1, S)_{P(1) \oplus P(n) \oplus \oplus(m-1)E_1}| = |Ext_\Lambda^1(\oplus mE_2, S)_{P(2) \oplus P(n+1) \oplus \oplus(m-1)E_2}| = q^m - 1,$$

$$g_{MN}^L = \frac{|Ext_\Lambda^1(MN)_L| |Aut(L)|}{|Aut(M)| |Aut(N)| |Hom|(M, N)|},$$

we obtain

$$g_{\oplus mE_1, S}^{P(1) \oplus P(n) \oplus \oplus(m-1)E_1} = g_{\oplus mE_2, S}^{P(2) \oplus P(n+1) \oplus \oplus(m-1)E_2} = q^{m-1}(q-1).$$

Applying Lemma 5.1, we get

$$g_{\oplus mE_1 \oplus \oplus rE_2, S}^{P(1) \oplus P(n) \oplus \oplus(m-1)E_1 \oplus \oplus rE_2} = q^{m-1}(q-1) |Hom_\Lambda(P(1) \oplus P(n), \oplus rE_2)|,$$

$$g_{\oplus mE_2 \oplus (\oplus rE_1), S}^{P(2) \oplus P(n+1) \oplus \oplus(m-1)E_2 \oplus \oplus rE_1} = q^{m-1}(q-1) |Hom_\Lambda(P(2) \oplus P(n+1), \oplus rE_1)|.$$

Moreover, by (A) and Lemma 16.13 in [KW], we have

$$V(S, \oplus mE_2 \oplus \oplus rE_1; P(1) \oplus P(n) \oplus (m-1)E_1 \oplus \oplus rE_2),$$

$$V(S, \oplus mE_2 \oplus \oplus rE_1; P(2) \oplus P(n+1) \oplus (m-1)E_2 \oplus \oplus rE_1)$$

have purity property.

Because of $Ext^1(\oplus mE_1, \oplus mE_1) = 0$, we get $\overline{\mathcal{O}}_{\oplus mE_1} = \mathbb{E}_{\dim \oplus mE_1}$.

So,

$$(p_3|_{\overline{\mathcal{O}}_{\oplus mE_1} \times \overline{\mathcal{O}}_S})^{-1}(P(1) \oplus P(n) \oplus (m-1)E_1) = P^{\dim Hom(S, P(1) \oplus P(n) \oplus (m-1)E_1) - 1}.$$

Thus,

$$(p_3|_{\overline{\mathcal{O}}_{\oplus mE_1} \times \overline{\mathcal{O}}_S})^{-1}(P(1) \oplus P(n) \oplus (m-1)E_1)$$

has purity property.

We now consider the cases $m \neq 0, r \neq 0$. By (A) and Mayer-Vietoris sequence, we only may consider the case $a = m, b = r$.

Consider the short exact sequence

$$(5.3) \quad 0 \longrightarrow S \longrightarrow P \longrightarrow \oplus mE_1 \oplus \oplus rE_2 \longrightarrow 0.$$

Applying $Hom(\oplus mE_1 \oplus \oplus rE_2,)$ to (5.3) we have

$$\begin{aligned}
0 &\longrightarrow \text{Hom}(\oplus mE_1 \oplus \oplus rE_2, S) \longrightarrow \text{Hom}(\oplus mE_1 \oplus \oplus rE_2, P) \\
&\longrightarrow \text{Hom}(\oplus mE_1 \oplus \oplus rE_2, \oplus mE_1 \oplus \oplus rE_2) \longrightarrow \text{Ext}^1(\oplus mE_1 \oplus \oplus rE_2, S) \\
&\longrightarrow \text{Ext}^1(\oplus mE_1 \oplus \oplus rE_2, P) \longrightarrow \text{Ext}^1(\oplus mE_1 \oplus \oplus rE_2, \oplus mE_1 \oplus \oplus rE_2) \longrightarrow 0
\end{aligned}$$

Since

$$\text{Hom}(\oplus mE_1 \oplus \oplus rE_2, S) = \text{Hom}(\oplus mE_1 \oplus \oplus rE_2, P) = 0,$$

we get

$$m^2 + r^2 \leq m + r.$$

Thus

$$m = r = 1.$$

We now have the short exact sequence

$$(5.4) \quad 0 \longrightarrow S \longrightarrow P \longrightarrow E_1 \oplus E_2 \longrightarrow 0.$$

If we also have the short exact sequence

$$(5.5) \quad 0 \longrightarrow S \longrightarrow P \xrightarrow{f} X \longrightarrow 0$$

where $X \not\cong E_1 \oplus E_2$.

Suppose P is indecomposable. Then by the projectivity of S , we obtain $\text{Hom}(P, S) = 0$. In addition, by $\text{End}(P) = k$, we have $\text{Hom}(X, P) = 0$.

Applying $\text{Hom}(\cdot, S)$ to (5.5), we have

$$\begin{aligned}
0 &\longrightarrow \text{Hom}(X, S) \longrightarrow \text{Hom}(P, S) \longrightarrow \text{Hom}(S, S) \\
&\longrightarrow \text{Ext}^1(X, S) \longrightarrow \text{Ext}^1(P, S) \longrightarrow \text{Ext}^1(S, S) \longrightarrow 0
\end{aligned}$$

Thus

$$\text{Hom}(X, S) = 0, \text{Ext}^1(X, S) = k^2.$$

Applying $\text{Hom}(X, \cdot)$ to (5.5) again, we have

$$\begin{aligned}
0 &\longrightarrow \text{Hom}(X, S) \longrightarrow \text{Hom}(X, P) \longrightarrow \text{Hom}(X, X) \\
&\xrightarrow{f^*} \text{Ext}^1(X, S) \longrightarrow \text{Ext}^1(X, P) \longrightarrow \text{Ext}^1(X, X) \longrightarrow 0
\end{aligned}$$

Because f^* is injective, we get $\dim_k \text{End}(X) \leq 2$. Thus $X \notin \overline{\mathcal{O}}_{E_1 \oplus E_2}$.

Suppose that P is decomposable. Since

$$\langle \delta, \underline{\dim} P \rangle = \langle \delta, \underline{\dim} S \rangle = -2,$$

we may assume that $P = P_1 \oplus P_2$ with P_1 and P_2 are pre-projective indecomposable. For convenience, we may assume that $n = 4$. Let $Y_i, i = 1, 2, 4, 5$ be pre-projective indecomposables of dimension vector $(01211), (10211), (11201), (11210)$, respectively. Then $P \cong P(1) \oplus Y_1$, or $P \cong P(2) \oplus Y_2$, or $P \cong P(4) \oplus Y_4$, or $P \cong P(5) \oplus Y_5$. Without loss of generality, we set $P = P(1) \oplus Y_1$.

Consider

$$(5.6) \quad 0 \longrightarrow S \longrightarrow P \longrightarrow E_1 \oplus E_2 \longrightarrow 0.$$

Since

$$\begin{aligned}
\dim \text{Hom}(P(1), E_1) &= \langle \underline{\dim} P(1), \underline{\dim} E_1 \rangle = 1, \dim \text{Hom}(P(1), E_2) = \langle \underline{\dim} P(1), \underline{\dim} E_2 \rangle = 0, \\
\dim \text{Hom}(Y_1, E_1) &= \langle \underline{\dim} Y_1, \underline{\dim} E_1 \rangle = 0, \dim \text{Hom}(Y_1, E_2) = \langle \underline{\dim} Y_1, \underline{\dim} E_2 \rangle = 1,
\end{aligned}$$

(5.6) turns into the following form

$$0 \longrightarrow S \longrightarrow P(1) \oplus Y_1 \begin{bmatrix} f_1 & 0 \\ 0 & f_2 \end{bmatrix} \longrightarrow E_1 \oplus E_2 \longrightarrow 0.$$

But $\dim \text{Im}(f_1) \leq 2 < 3 = \dim E_1$. This gives a contradiction. Thus P must be indecomposable.

We first point out that P is the extension of S by $E_1 \oplus E_2$. In fact, $P \cong M(x)$, and $E_1 \oplus E_2 \cong M(y)$, where $x_{13} = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$, $x_{23} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $x_{43} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, $x_{53} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$; $y_{13} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $y_{23} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$, $y_{43} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $y_{53} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. It is easy to see that $(f_i)_i$ is an epimorphism from P to $E'_1 \oplus E'_2$, here $f_1 = f_5 = -1$, $f_2 = f_4 = 1$, $f_3 = \begin{bmatrix} 0 & 1 & -1 \\ 1 & 0 & -1 \end{bmatrix}$.

From the discussions above, we obtain

$$(p_3|_{\overline{\mathcal{O}}_{E_1 \oplus E_2} \times \mathcal{O}_S})^{-1}(P) = V(S, E_1 \oplus E_2; P).$$

Let X be the set of epimorphisms from P to $E_1 \oplus E_2$. Then

$$V(S, E_1 \oplus E_2; P) = X / \text{Aut}(E_1 \oplus E_2).$$

Here the action of $\text{Aut}_\Lambda(E_1 \oplus E_2)$ on X is defined by

$$h \circ f = hf.$$

Since

$$\begin{aligned} \text{Hom}_\Lambda(P, E_1 \oplus E_2) &= \{(f_i)_i | f_1 = a, f_2 = b, f_4 = -a, f_5 = -b, f_3 = \begin{bmatrix} 0 & -a & -a \\ b & 0 & -b \end{bmatrix}, a, b \in k\}, \\ \text{End}_\Lambda(E_1 \oplus E_2) &= \{(h_i)_i | h_1 = c, h_2 = d, h_4 = c, h_5 = d, h_3 = \begin{bmatrix} c & 0 \\ 0 & d \end{bmatrix}, c, d \in k\}, \end{aligned}$$

we have

$$\begin{aligned} X &= \{(f_i)_i | (f_i)_i \in \text{Hom}_\Lambda(P, E_1 \oplus E_2), a \neq 0, b \neq 0\}, \\ \text{Aut}_\Lambda(E_1 \oplus E_2) &= \{(h_i)_i | (h_i)_i \in \text{End}_\Lambda(E_1 \oplus E_2), c \neq 0, d \neq 0\}. \end{aligned}$$

Then $X / \text{Aut}(E_1 \oplus E_2)$ has only one point correspond to the orbit of $(1, 1, \begin{bmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \end{bmatrix}, -1, -1)$, that is,

$$V(S, E_1 \oplus E_2; P) = \{1 \text{ point}\}.$$

Thus $p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2} \times \mathcal{O}_S})^{-1}(P)$ has purity property.

Case 3 : $Q = \tilde{E}_6$.


We only consider $\mathcal{T}_1$ in Fig 4.2.3. Other cases can be proved similarly. Let $E_i, i = 1, 2$ be simple objects in the full subcategory $\mathcal{T}_1$ of $\Lambda - \text{mod}$ corresponding to the dimension vectors listed in Fig 4.2.3.

Since

$$\dim \text{Ext}^1(E_1, S) = -\langle \underline{\dim} E_1, \underline{\dim} S \rangle = 1, \text{ and } \dim \text{Ext}^1(E_2, S) = -\langle \underline{\dim} E_2, \underline{\dim} S \rangle = 2,$$

we have

$$\dim \text{Ext}^1(\oplus m E_1, S) = m, \dim \text{Ext}^1(\oplus m E_2, S) = 2m.$$



where $P(3)=S$.

Fig.5.2

Let Y, N be any Λ -modules, and let $N_1 = S_6, N_2 = S_2, N_3 = S_4$. We denote by W_N the set of maps $w : S \rightarrow Y$ such that either $w = 0$ or $\text{Hom}_\Lambda(N, \text{Cok} w) \neq 0$, and denote by W_t the set of maps $w : S \rightarrow Y$ which factor through $P(t)$. Suppose $\text{Hom}_\Lambda(S, Y) = k^{n+1}$, then the coordinate ring of $\text{Hom}_\Lambda(S, Y)$ is $k[T_0, \dots, T_n]$.

The action of $k^* = \text{Aut}(S)$ on $\text{Hom}_\Lambda(S, Y) = k^{n+1}$ is defined by

$$a \circ (w_0, \dots, w_n) = (w_0 a, \dots, w_n a).$$

When $Y = X$, we have $\text{Hom}_\Lambda(S, X) = k^2$, and the coordinate ring of $\text{Hom}_\Lambda(S, X)$ is $k[T_0, T_1]$. Let r_{N_j} be the defining relations of W_{N_j} , then $r_{N_1} = T_1, r_{N_2} = T_0, r_{N_3} = T_0 - T_1$, that is, $W_{N_j} = \mathcal{L}(r_{N_j})$ for $j = 1, 2, 3$.

Thanks to the Corollary on page 166 in [R3], we get

$$\mathbf{Z}_{E'_2, S, X} = (\mathbb{A}^2 - \cup_{j=1}^3 \mathcal{L}(r_{N_j})) / k^* = P^1 - \{[1 : 0], [0 : 1], [1 : 1]\}.$$

Since

$$\dim_k \text{Hom}_\Lambda(S, E_2) = \langle \dim S, \dim E_2 \rangle = 1,$$

we have

$$\text{Hom}_\Lambda(S, X \bigoplus \oplus(m-1)E_2) = k^{m+1}.$$

For any $0 \neq w \in \text{Hom}_\Lambda(S, X \bigoplus \oplus(m-1)E_2)$, suppose

$$\text{Cok} w = a_1 Y_1 \oplus a_1 Y_1 \oplus a_2 Y_2 \oplus a_3 Y_3 \oplus (m-i)E_2 \oplus b_1 N_1 \oplus b_2 N_2 \oplus b_3 N_3.$$

then we have

$$(5.7) \quad \begin{cases} a_1 + a_2 + a_3 &= i \\ a_2 + a_3 + b_1 &= i \\ a_3 + a_1 + b_2 &= i \\ a_1 + a_2 + b_3 &= i. \end{cases}$$

Then $a_j = b_j$, for all j , and $b_j = 0, a_j = 0$ if $\text{Hom}_\Lambda(N_j, \text{Cok} w) = 0$ for all j , that is, $\text{Cok} w \cong \oplus m E_2$. From Fig.5.2, we, again, have $r_{N_1} = T_1, r_{N_2} = T_0, r_{N_3} = T_0 - T_1$.

According to 7.2 in [R3], we obtain

$$(5.8) \quad \mathbf{Z}_{\oplus m E_2, S, X \bigoplus \oplus(m-1)E_2} = (\mathbb{A}^{m+1} - \cup_{j=1}^3 \mathcal{L}(r_{N_j})) / k^*.$$

As to the triple modules $(\oplus m E_2, S, P(6) \oplus Y_1 \bigoplus \oplus(m-1)E_2)$, by comparing dimension vectors of middle term and both ends in exact sequence, similar to 5.7, we obtain that N_i is still the test module of a pair modules $(P(6) \oplus Y_1 \bigoplus \oplus(m-1)E_2, S)$.

So, we also have

$$\mathbf{Z}_{\oplus m E'_2, S, P(6) \oplus Y_1 \bigoplus \oplus(m-1)E'_2} = (\mathbb{A}^{m+1} - \cup_{j=1}^3 \mathcal{L}(r_{N_j})) / k^*.$$

Thus

$$\mathbf{Z}_{\oplus m E_2, S, X \bigoplus \oplus(m-1)E_2}, \mathbf{Z}_{\oplus m E'_2, S, P(6) \oplus Y_1 \bigoplus \oplus(m-1)E'_2}$$

have purity property.

Now, from Fig.5.2, we have

$$\begin{aligned} & p_3|_{\overline{\mathcal{O}}_{\oplus mE_2} \times \mathcal{O}_S}^{-1}(X \oplus \oplus(m-1)E_2) \\ &= \mathbf{Z}_X \oplus_{(m-1)E_2, \oplus mE_2, S \cup \cup_{i=1}^3 \mathcal{L}(rN_i)} \\ &= (Hom(S, X \oplus \oplus(m-1)E_2) - \{0\})/k^* = P^{dim_k Hom(S, X \oplus \oplus(m-1)E_2)-1}. \end{aligned}$$

Similarly,

$$\begin{aligned} & p_3|_{\overline{\mathcal{O}}_{\oplus mE_1} \times \mathcal{O}_S}^{-1}(P(1) \oplus P(5) \oplus P(7) \oplus (m-1)E_1) \\ &= p_3^{-1}(\tau^{-1}(P(1) \oplus P(5) \oplus P(7)) \oplus \oplus(m-1)E_2) \\ &= p_3^{-1}((Y_1 \oplus Y_2 \oplus Y_3) \oplus \oplus(m-1)E_2) \\ &= P^{dim Hom(X, (Y_1 \oplus Y_2 \oplus Y_3) \oplus \oplus(m-1)E_2)-1}. \end{aligned}$$

Thus the statement is true if $m = 0$ or $r = 0$.

We now assume that $m \neq 0, r \neq 0$. By (A) and Mayer-Vietoris sequence, we also only need to consider the case $a = m, b = r$

Consider the short exact sequence

$$(5.9) \quad 0 \longrightarrow S \longrightarrow P \xrightarrow{g} \oplus mE_1 \bigoplus rE_2 \longrightarrow 0.$$

Applying $Hom(\oplus mE_1 \bigoplus \oplus rE_2,)$ to (5.9), we obtain

$$\begin{aligned} 0 \longrightarrow Hom(\oplus mE_1 \bigoplus \oplus rE_2, S) \longrightarrow Hom(\oplus mE_1 \bigoplus \oplus rE_2, P) \longrightarrow Hom(\oplus mE_1 \bigoplus \oplus rE_2, \oplus mE_1 \bigoplus \oplus rE_2) \\ \xrightarrow{g^*} Ext^1(\oplus mE_1 \bigoplus \oplus rE_2, S) \longrightarrow Ext^1(\oplus mE_1 \bigoplus \oplus rE_2, P) \longrightarrow Ext^1(\oplus mE_1 \bigoplus \oplus rE_2, S) \longrightarrow 0 \end{aligned}$$

Because g^* is an injective, we get $m^2 + r^2 \leq m + 2r$. Thus $m = 1; r = 1, 2$.

When $m = r = 1$, we consider the short exact sequence

$$(5.10) \quad 0 \longrightarrow S \longrightarrow P \longrightarrow E_1 \oplus E_2 \longrightarrow 0.$$

Since $\langle \delta, \underline{\dim} S \rangle = \langle \delta, \underline{\dim} S \rangle = -3$, P is an indecomposable object, or the sum of two (or three) indecomposable objects.

Suppose that $P \cong P_1 \oplus P_2 \oplus P_3$ with P_i to be indecomposable for $i = 1, 2, 3$. By the AR -quiver of $\tilde{E}_6$, we have

$$P \cong P(1) \oplus \tau^{-1}(P(1)) \oplus \tau^{-2}(P(1)), \text{ or } P(5) \oplus \tau^{-1}(P(5)) \oplus \tau^{-2}(P(5)), \text{ or } P(7) \oplus \tau^{-1}(P(7)) \oplus \tau^{-2}(P(7)).$$

Without loss of generality, we may assume that $P \cong P(1) \oplus \tau^{-1}(P(1)) \oplus \tau^{-2}(P(1))$. Because there is not epimorphism from $\tau^{-1}(P(1))$ to E_2 , by $Hom(P(1) \oplus \tau^{-2}(P(1)), E_2) = 0$, there is not exact sequence

$$0 \longrightarrow S \longrightarrow P(1) \oplus \tau^{-1}(P_1) \oplus \tau^{-2}(P_1) \longrightarrow E_1 \oplus E_2 \longrightarrow 0$$

Suppose that P is an indecomposable extension of S by $E_1 \oplus E_2$. Let X be a module in $\overline{\mathcal{O}}_{E_1 \oplus E_2} \setminus \mathcal{O}_{E_1 \oplus E_2}$, and we have the following the short exact sequence

$$(5.11) \quad 0 \longrightarrow S \longrightarrow P \longrightarrow X \longrightarrow 0.$$

Applying $Hom(, S)$ (resp. $Hom(X,)$) to (5.11), we have

$$\begin{aligned} 0 \longrightarrow Hom(X, S) \longrightarrow Hom(P, S) \longrightarrow Hom(S, S) \\ \longrightarrow Ext^1(X, S) \longrightarrow Ext^1(P, S) \longrightarrow Ext^1(S, S) \longrightarrow 0 \end{aligned}$$

and

$$\begin{aligned} 0 \longrightarrow Hom(X, S) \longrightarrow Hom(X, P) \longrightarrow Hom(X, X) \\ \longrightarrow Ext^1(X, S) \longrightarrow Ext^1(X, P) \longrightarrow Ext^1(X, X) \longrightarrow 0. \end{aligned}$$

Now, it follows from $\text{Hom}(P, S) = 0$ and $\text{Hom}(X, P) = 0$ that

$$\text{Hom}(X, S) = 0, \text{Ext}(X, S) = k^3, \dim \text{End}(X) \leq 3.$$

According to the assumption on X , we obtain $\dim \text{End}(X) = 3$.

If $X \cong X_1 \oplus X_2$ and $X_i \in \mathcal{T}_1, i = 1, 2$, then we have $X \cong E_1 \oplus E_2$, a contradiction.

If $X \cong X_1 \oplus X_2$ and $X_1 \in \mathcal{P}_{\text{prep}}, X_2 \in \mathcal{T}_1$ (resp. $X_1 \in \mathcal{T}_1, X_2 \in \mathcal{P}_{\text{prei}}$), by Proposition 4.1.2, then $E_1 \oplus E_2$ is an extension X_1 by X_2 . This result contradicts to the theory of representations of tame quivers.

If $X \cong X_1 \oplus X_2$ and $X_1 \in \mathcal{P}_{\text{prep}}, X_2 \in \mathcal{P}_{\text{prei}}$. Then $\dim \text{Hom}(X_1, X_2) = 1$. Suppose $X_1 = \tau^{-a}P(j)$, then we have

$$1 = \dim \text{Hom}(\tau^{-a}P(j), X_2) = \dim \text{Hom}(P(j), \tau^a X_2).$$

It implies that $X_1 = \tau^{-a}P(2), \tau^{-a}P(4)$, or $\tau^{-a}P(6)$. According to the AR -quiver of $\tilde{E}_6$, we have $a = 2$ and $X_2 = I(2), I(4)$, or $I(6)$.

If $X \cong X_1 \oplus X_2 \oplus X_3$, we have $\text{Hom}(X_i, X_j) = 0$ fore all $i \neq j$. Suppose that there is one regular term in $X_i, i = 1, 2, 3$, and the other two terms are pre-projective or pre-injective, then, similar to the above discussions, we get $X \notin \overline{\mathcal{O}}_{E_1 \oplus E_2}$. Suppose X_3 (resp. X_1) is pre-injective (resp. pre-projective), and the other two terms are pre-projective (resp. pre-injective). By $X \in \overline{\mathcal{O}}_{E_1 \oplus E_2}$ and $\dim \text{End}(X) = 3, \dim \text{End}(E_1 \oplus E_2) = 2$, we have

$$(5.12) \quad 0 \longrightarrow X_1 \oplus X_2 \longrightarrow E_1 \oplus E_2 \longrightarrow X_3 \longrightarrow 0.$$

Applying $\text{Hom}(_, X_3)$ to (5.12), we obtain

$$\begin{aligned} 0 &\longrightarrow \text{Hom}(X_3, X_3) \longrightarrow \text{Hom}(E_1 \oplus E_2, X_3) \longrightarrow \text{Hom}(X_1 \oplus X_2, X_3) \\ &\longrightarrow \text{Ext}^1(X_3, X_3) \longrightarrow \text{Ext}^1(E_1 \oplus E_2, X_3) \longrightarrow \text{Ext}^1(X_1 \oplus X_2, X_3) \longrightarrow 0. \end{aligned}$$

Because of $\text{Hom}(X_1 \oplus X_2, X_3) = 0$, we can deduce $\dim \text{Hom}(E_1 \oplus E_2, X_3) = 1$.

Applying $\text{Hom}(E_1 \oplus E_2, _)$ to (5.12), we also obtain $\dim \text{Hom}(E_1 \oplus E_2, X_3) \geq 2$ by $\text{Hom}(E_1 \oplus E_2, X_1 \oplus X_2) = 0$. This gives a contradiction.

Let X_1 (resp. X_2, X_3) be the pre-projective (resp. regular, pre-injective) indecomposable module so that $X_2 \cong E_1$ or $X_2 \cong E_2$. Suppose $X_2 \cong E_1$, then it follows from (5.12) that

$$0 \longrightarrow X_1 \longrightarrow E_2 \longrightarrow X_3 \longrightarrow 0.$$

Thus $X_1 \cong \tau^{-1}P(1)$, or $X_1 \cong \tau^{-1}P(5)$, or $X_1 \cong \tau^{-1}P(7)$.

Suppose $X_2 \cong E_2$, then it follows from (5.12) that

$$0 \longrightarrow X_1 \longrightarrow E_1 \longrightarrow X_3 \longrightarrow 0.$$

Thus $X_3 \cong I(1)$, or $X_3 \cong I(5)$, or $X_3 \cong I(7)$.

We, thus, only need to consider (I) and (II).

(I) $X \cong X_1 \oplus X_2, X_1 \in \mathcal{P}_{\text{prep}}, X_2 \in \mathcal{P}_{\text{prei}}$, and $\text{Hom}(X_1, X_2) = 1$. Then, $X \cong I(2) \oplus \tau^{-2}P(2)$ (resp. $I(4) \oplus \tau^{-2}P(4), I(6) \oplus \tau^{-2}P(6)$).

Fix an exact sequence as follows

$$0 \longrightarrow P(1) \xrightarrow{p} P \xrightarrow{q} \tau^{-2}P(2) \longrightarrow 0.$$

Since

$$\dim \text{Hom}(P(1), I(2)) = \dim \text{Hom}(S, P(1)) = 1,$$

we choose $u_0 \in \text{Hom}(S, P(1))$ (resp. $v_0 \in \text{Hom}(P(1), I(2))$) such that $\{u_0\}$ (resp. $\{v_0\}$) is the basis of $\text{Hom}(S, P(1))$ (resp. $\text{Hom}(P(1), I(2))$).

For any $(u, v) \in W(S, P(1); I(2))$, we have following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & S & \xrightarrow{u} & P(1) & \xrightarrow{v} & I(2) \longrightarrow 0 \\ & & \parallel & & p \downarrow & & \downarrow \\ 0 & \longrightarrow & S & \xrightarrow{pu} & P & \xrightarrow{\pi} & Cok(pu) \longrightarrow 0. \end{array}$$

Based on the Snake Lemma, we get $Cok(pu) \cong I(2) \oplus \tau^{-2}P(2)$. Thus $(pu, w\pi) \in W(S, I(2) \oplus \tau^{-2}P(2); P)$ by $Hom(I(2), P) = 0$.

Conversely, let $(f, g) \in W(S, I(2) \oplus \tau^{-2}P(2); P)$, by the projectivity of $P(1)$, there exists morphisms $u = au_0, p' = bp$ such that we also have the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & S & \xrightarrow{u} & P(1) & \xrightarrow{v_0} & I(2) \longrightarrow 0 \\ & & \parallel & & p' \downarrow & & \left[\begin{smallmatrix} 1 \\ 0 \end{smallmatrix} \right] \downarrow \\ 0 & \longrightarrow & S & \xrightarrow{f} & P & \xrightarrow{g} & I(2) \oplus \tau^{-2}P(2) \longrightarrow 0. \end{array}$$

Thus $(bu, b^{-1}v_0) \in W(S, I(2); P(1))$. We deduce that

$$V(S, I(2) \oplus \tau^{-2}P(2); P) = V(S, I(2); P(1)) = \{1 \text{ point}\}.$$

(II) $X \cong X_1 \oplus X_2 \oplus X_3, X_1 \in \mathcal{P}_{prep}, X_2 \in \mathcal{P}_{regular}, X_3 \in \mathcal{P}_{prei}$, and $Hom(X_i, X_j) = 0, i \neq j$. Thus, $X \cong \tau^{-1}P(1) \oplus E_1 \oplus \tau I(1)$ (resp. $\tau^{-1}P(5) \oplus E_1 \oplus \tau I(5), \tau^{-1}P(7) \oplus E_1 \oplus \tau I(7)$) or $X \cong \tau^{-2}P(1) \oplus E_2 \oplus I(1)$ (resp. $X \cong \tau^{-2}P(5) \oplus E_2 \oplus I(5), X \cong \tau^{-2}P(7) \oplus E_2 \oplus I(7)$).

Since

$$Hom(P, \tau^{-1}P(1)) = Hom(P, \tau^{-1}P(5)) = Hom(P, \tau^{-1}P(7)) = 0$$

and

$$Ext^1(I(1), S) = Ext^1(I(5), S) = Ext^1(I(7), S) = 0,$$

there is not exact sequence of form (5.11).

From the discussions above, we obtain

$$\begin{aligned} p_3|_{\overline{\mathcal{O}}_{E_1 \oplus E_2} \times \mathcal{O}_S}^{-1}(P) \\ = V(S, E_1 \oplus E_2; P) \cup V(S, I(2) \oplus \tau^{-2}P(2); P) \\ \cup V(S, \tau^{-2}P(4) \oplus I(4); P) \cup V(S, \tau^{-2}P(6) \oplus I(6); P). \end{aligned}$$

We first point out that P is the extension of S by $E_1 \oplus E_2$. In fact, $P \cong M(x)$, and $E_1 \oplus E_2 \cong$

$$M(y), \text{ where } x_{12} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, x_{23} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, x_{43} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}, x_{54} = \begin{bmatrix} 0 \\ 1 \end{bmatrix},$$

$$x_{63} = \begin{bmatrix} 0 & 1 \\ 1 & 1 \\ 1 & 1 \\ 0 & 1 \end{bmatrix}, x_{76} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}; y_{12} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, y_{23} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}, y_{43} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}, y_{54} = \begin{bmatrix} 1 \\ 0 \end{bmatrix},$$

$$y_{63} = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}, y_{76} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \text{ It is easy to see that } (f_i)_i \text{ is an epimorphism from } P \text{ to } E_1 \oplus E_2,$$

where $f_1 = 1, f_2 = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, f_3 = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & -1 & 0 & 0 \end{bmatrix}, f_4 = \begin{bmatrix} 1 & 1 \\ -1 & 0 \end{bmatrix}, f_5 = 1,$

$$f_6 = \begin{bmatrix} 1 & 2 \\ -1 & 0 \end{bmatrix}, f_7 = 2.$$

Let X be the set of epimorphisms from P to $E_1 \oplus E_2$. Then

$$V(S, E_1 \oplus E_2; P) = X / \text{Aut}(E_1 \oplus E_2).$$

Here the action of $\text{Aut}_\Lambda(E_1 \oplus E_2)$ on X is defined by

$$h \circ f = hf.$$

Since

$$\text{Hom}_\Lambda(P, E_1 \oplus E_2)$$

$$= \{(f_i)_i | f_1 = a, f_2 = \begin{bmatrix} b_1 & a \\ b_2 & 0 \end{bmatrix}, f_3 = \begin{bmatrix} b_1 & 0 & a & 0 \\ 0 & a & 0 & b_1 \\ b_2 & -b_2 & 0 & 0 \end{bmatrix}, f_4 = \begin{bmatrix} a & b_1 \\ -b_2 & 0 \end{bmatrix}, f_5 = b_1,$$

$$f_6 = \begin{bmatrix} a & a + b_1 \\ -b_2 & 0 \end{bmatrix}, f_7 = a + b_1; a, b_1, b_2 \in k\},$$

$$\text{End}_\Lambda(E_1 \oplus E_2)$$

$$= \{(h_i)_i | h_1 = c, h_2 = \begin{bmatrix} c & 0 \\ 0 & d \end{bmatrix}, h_3 = \begin{bmatrix} c & 0 & 0 \\ 0 & c & 0 \\ 0 & 0 & d \end{bmatrix}, h_4 = \begin{bmatrix} c & 0 \\ 0 & d \end{bmatrix}, h_5 = c,$$

$$h_6 = \begin{bmatrix} c & 0 \\ 0 & d \end{bmatrix}, h_7 = c; c, d \in k\}\},$$

Then we have

$$X = \{(f_i)_i | (f_i)_i \in \text{Hom}_\Lambda(P, E_1 \oplus E_2), a \neq 0, b_1 \neq 0, b_2 \neq 0, a + b_1 \neq 0\},$$

$$\text{Aut}_\Lambda(E_1 \oplus E_2) = \{(h_i)_i | (h_i)_i \in \text{End}_\Lambda(E_1 \oplus E_2), c \neq 0, d \neq 0\}.$$

Hence $X / \text{Aut}(E_1 \oplus E_2)$ have points corresponds to orbits of $(f_1 = 1, f_2 = \begin{bmatrix} b_1/a & 1 \\ 1 & 0 \end{bmatrix},$

$$f_3 = \begin{bmatrix} b_1/a & 0 & 1 & 0 \\ 0 & 1 & 0 & b_1/a \\ 1 & -1 & 0 & 0 \end{bmatrix}, f_4 = \begin{bmatrix} 1 & b_1/a \\ -1 & 0 \end{bmatrix}, f_5 = b_1/a, f_6 = \begin{bmatrix} 1 & 1 + b_1/a \\ -1 & 0 \end{bmatrix},$$

$f_7 = 1 + b_1/a; b_1/a \neq 0, 1$, that is,

$$V(S, E_1 \oplus E_2; P) = \mathbb{A}^1 \setminus \{2 \text{ point}\}.$$

By Lemma 16.13 in [KW], we have $V(S, E_1 \oplus E_2; P)$ has purity property.

Thereby $p_3|_{\overline{\mathcal{O}}_{E_1 \oplus E_2} \times \mathcal{O}_S}^{-1}(P)$ has purity property.

Suppose that $P \cong P_1 \oplus P_2$ with P_i is indecomposable for $i = 1, 2$. By the AR -quiver of $\tilde{E}_6$, we have

$$P \cong P(1) \oplus \tau^{-2}(P(2)), \text{ or } P(5) \oplus \tau^{-2}(P(4)), \text{ or } P(7) \oplus \tau^{-2}(P(6));$$

$$P \cong \tau^{-2}P(1) \oplus \tau^{-1}(P(2)), \text{ or } \tau^{-2}P(5) \oplus \tau^{-1}(P(4)), \text{ or } \tau^{-2}P(7) \oplus \tau^{-1}(P(6)).$$

Without loss of generality, we assume that $P \cong P(1) \oplus \tau^{-2}(P(2))$. We claim that P is the nonzero extension of S by $E_1 \oplus E_2$.

Let $x_{12} = 0, x_{23} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, x_{43} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}, x_{54} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, x_{63} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}, x_{76} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$. We then have $\tau^{-2}(P(2)) = M(x)$.

It is easy to see that $f = (f_i)_i$ (resp. $\vartheta = (\vartheta_i)_i$) is the epimorphism from $\tau^{-2}(P(2)) \oplus P(1)$ (resp. $\tau^{-2}(P(2))$) to $E_1 \oplus E_2$ (resp. E_2), where $f_1 = 1, f_2 = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, f_3 = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & -1 & 0 & 0 \end{bmatrix}, f_4 = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, f_5 = 1, f_6 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, f_7 = 1$ (resp. $\vartheta_1 = 0, \vartheta_2 = 2, \vartheta_3 = \begin{bmatrix} 1 & 1 & 0 \end{bmatrix}, \vartheta_4 = \begin{bmatrix} 1 & 0 \end{bmatrix}, \vartheta_5 = 0, \vartheta_6 = \begin{bmatrix} 1 & 0 \end{bmatrix}, \vartheta_7 = 0$.) So we get that f is the epimorphism from $P(1) \oplus \tau^{-1}(P(2))$ to $E_1 \oplus E_2$. The claim is proven.

Suppose that there is X with $E_1 \oplus E_2 \not\cong X$ and $X \in \overline{\mathcal{O}}_{E_1 \oplus E_2}$ such that there is the short exact sequence

$$(5.13) \quad 0 \longrightarrow S \longrightarrow P(1) \oplus \tau^{-2}(P(2)) \longrightarrow X \longrightarrow 0.$$

If $\text{Hom}(X, P(1) \oplus \tau^{-2}(P(2))) = 0$, then we have $\dim \text{End}(X) = 3$. Similar to the case of the indecomposable P , we deduce that there does not exist X such that (5.13) holds.

Let now $\text{Hom}(X, P(1) \oplus \tau^{-2}(P(2))) \neq 0$. There is an indecomposable summand $\tau^{-a}P(j)$ of X such that $\text{Hom}(\tau^{-a}P(j), P(1) \oplus \tau^{-2}(P(2))) \neq 0$. Set $X_1 = \tau^{-a}P(j)$ and $X = X_1 \oplus X_2$.

If $X_1 = P(1)$, by $\text{Hom}(P(1), X_2) = 0$, we have

$$0 \longrightarrow S \longrightarrow \tau^{-2}(P(2)) \longrightarrow X_2 \longrightarrow 0,$$

and

$$p_3|_{\overline{\mathcal{O}}_{X_2 \times \mathcal{O}_S}}^{-1}(\tau^{-2}(P(2))) \cong p_3|_{\overline{\mathcal{O}}_{P(1) \oplus X_2 \times \mathcal{O}_S}}^{-1}(P(1) \oplus \tau^{-2}(P(2))).$$

If $X_1 \not\cong P(1)$, X_1 must be non-projective. According to the AR -quiver of $\tilde{E}_6$, we have $X_1 = \tau^{-a}P(j); j = 1, 2, 3, 4, 5, 6, 7; a = 1, 2$.

Suppose $a \neq 2, j \neq 2$. For any $f \in \text{Hom}(P(1), \tau^{-a}P(j))$, we know that f is not an epimorphism. Thus $a = 2, j = 2$, and $\underline{\dim} X_2 = \underline{\dim} I(2)$. Because of $\text{Ext}^1(I(1), S) = 0$, it follows from (5.13) that $X_2 = I(2)$, that is, we get the following short exact sequence

$$0 \longrightarrow S \longrightarrow P(1) \oplus \tau^{-2}P(2) \longrightarrow I(2) \oplus \tau^{-2}(P(2)) \longrightarrow 0.$$

From the discussions above, we obtain

$$\begin{aligned} & p_3|_{\overline{\mathcal{O}}_{E_1 \oplus E_2 \times \mathcal{O}_S}}^{-1}(P(1) \oplus \tau^{-2}P(2)) \\ &= V(S, E_1 \oplus E_2; P(1) \oplus \tau^{-2}P(2)) \cup V(S, I(2) \oplus \tau^{-2}P(2); P(1) \oplus \tau^{-2}P(2)) \\ & \quad \cup_{[X_2]} p_3|_{\overline{\mathcal{O}}_{P(1) \oplus X_2 \times \mathcal{O}_S}}^{-1}(P(1) \oplus \tau^{-2}(P(2))), \end{aligned}$$

where $\cup$ denote the disjoint union.

Obviously,

$$V(S, I(2) \oplus \tau^{-2}P(2); P(1) \oplus \tau^{-2}P(2)) = \{1 \text{ point}\}.$$

Moreover,

$$p_3|_{\overline{\mathcal{O}}_{P(1) \oplus X_2 \times \mathcal{O}_S}}^{-1}(P(1) \oplus \tau^{-2}(P(2)))$$

has purity property by [L4].

Let $z_{12} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$, $z_{23} = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}$, $z_{43} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$, $z_{54} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $z_{63} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$,
 $z_{76} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$. Then $\tau^{-2}P(2) \oplus P(1) \cong M(z)$.

Since

$$\begin{aligned} & Hom_{\Lambda}(\tau^{-2}P(2) \oplus P(1), E_1 \oplus E_2) \\ &= \{(f_i)_i | f_1 = a, f_2 = \begin{bmatrix} b_1 & a \\ b_2 & 0 \end{bmatrix}, f_3 = \begin{bmatrix} b_1 & 0 & 0 & a \\ b_1 & -b_1 & 0 & 0 \\ 0 & 0 & b_2 & 0 \end{bmatrix}, f_4 = \begin{bmatrix} -b_1 & 0 \\ 0 & b_2 \end{bmatrix}, f_5 = -b_1, \\ & f_6 = \begin{bmatrix} b_1 & 0 \\ 0 & b_2 \end{bmatrix}, f_7 = b_1; a, b_1, b_2 \in k\}, \end{aligned}$$

Let X be the set of epimorphisms from $\tau^{-2}P(2) \oplus P(1)$ to $E_1 \oplus E_2$. Then we have

$$X = \{(f_i)_i | (f_i)_i \in Hom_{\Lambda}(P, E_1 \oplus E_2), a \neq 0, b_1 \neq 0, b_2 \neq 0\}.$$

Hence $X/Aut(E_1 \oplus E_2)$ have points correspond to orbits of $(f_1 = 1, f_2 = \begin{bmatrix} b_1/a & 1 \\ 1 & 0 \end{bmatrix})$,
 $f_3 = \begin{bmatrix} b_1/a & 0 & 0 & 1 \\ b_1/a & -b_1/a & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$, $f_4 = \begin{bmatrix} -b_1/a & 0 \\ 0 & 1 \end{bmatrix}$, $f_5 = -b_1/a$, $f_6 = \begin{bmatrix} b_1/a & 0 \\ 0 & 1 \end{bmatrix}$,
 $f_7 = b_1/a$; $b_1/a \neq 0$, that is,

$$V(S, E_1 \oplus E_2; P) = \mathbb{A}^1 \setminus \{1 \text{ point}\}.$$

By Lemma 16.13 in [KW], we have $V(S, E_1 \oplus E_2; P)$ has purity property.

Thus $p_3|_{\overline{\mathcal{O}}_{E_1 \oplus E_2} \times \mathcal{O}_S}^{-1}(P(1) \oplus \tau^{-2}P(2))$ has purity property.

When $m = 1, r = 2$, let P be the extension of S by $E_1 \oplus 2E_2$. Since $\langle \delta, \underline{\dim} P \rangle = \langle \delta, \underline{\dim} S \rangle = -3$, we can deduce that P has 3 indecomposable direct summands at most.

If P is also the extension of S by X with $X \in \overline{\mathcal{O}}_{E_1 \oplus 2E_2} \setminus \mathcal{O}_{E_1 \oplus 2E_2}$, that is,

$$(5.14) \quad 0 \longrightarrow S \longrightarrow P \longrightarrow X \longrightarrow 0.$$

Applying $Hom(\cdot, S)$ and $Hom(X, \cdot)$ to (5.14), we get $\dim End(X) \leq 5 + \dim Hom(X, P)$.

Suppose P is an indecomposable. It follows from $Hom(X, P) = 0$ that $\dim End(X) \leq 5$. But $\dim End(E_1 \oplus 2E_2) = 5$, so there does not exist $X \in \overline{\mathcal{O}}_{E_1 \oplus 2E_2} \setminus \mathcal{O}_{E_1 \oplus 2E_2}$ such that (5.14) holds. Then, similar to $m = r = 1$ case,

$$p_3|_{\overline{\mathcal{O}}_{E_1 \oplus 2E_2} \times \mathcal{O}_S}^{-1}(P) = V(S, E_1 \oplus 2E_2; P)$$

has purity property.

Suppose that P can be decomposed into two indecomposable objects, say, $P \cong P_1 \oplus P_2$ such that

$$\begin{aligned} P &\cong \tau^{-1}P(1) \oplus \tau^{-3}P(2) \text{ or } \tau^{-1}P(5) \oplus \tau^{-3}P(4) \text{ or } \tau^{-1}P(7) \oplus \tau^{-3}P(6) \\ P &\cong \tau^{-2}P(2) \oplus \tau^{-3}P(1) \text{ or } \tau^{-2}P(4) \oplus \tau^{-3}P(5) \text{ or } \tau^{-1}P(6) \oplus \tau^{-3}P(7). \end{aligned}$$

Without loss generality, we assume that $P \cong \tau^{-1}P(1) \oplus \tau^{-3}P(2)$.

If $Hom(X, P) = 0$, it follows from $\dim End(X) \leq 5$ that $X \notin \overline{\mathcal{O}}_{E_1 \oplus 2E_2}$.

If $\text{Hom}(X, P) \neq 0$, similar to the case of $m = r = 1$, we get $X \cong \tau^{-1}P(1) \oplus X_2$ and $\text{Hom}(X_2, P) = 0$ or $X \cong X_1 \oplus \tau^{-3}P(2)$.

When $X \cong X_1 \oplus \tau^{-3}P(2)$, we have $X_1 = \tau I(5) \oplus \tau I(7)$ and

$$g_{\tau I(5) \oplus \tau I(7), S}^{\tau^{-1}P(1)} = g_{\tau I(5) \oplus \tau I(7) \oplus \tau^{-3}P(2), S}^{\tau^{-1}P(1) \oplus \tau^{-3}P(2)} = 1.$$

That is,

$$V(S, \tau I(5) \oplus \tau I(7) \bigoplus \tau^{-3}P(2); \tau^{-1}P(1) \oplus \tau^{-3}P(2)) = \{1 \text{ point}\}.$$

When $X \cong \tau^{-1}P(1) \oplus X_2$. We then have $\dim \text{End}(X) = 5 + \dim \text{Hom}(X, P) = 6$. Thus X_2 must be decomposable, and $\dim \text{End}(X_2) = 4$.

Suppose $X_2 \cong X_{21} \oplus X_{22}$ and $X_{21} \in \mathcal{P}_{\text{prep}}, X_{22} \in \mathcal{P}_{\text{prei}}$. It follows from $\dim \text{End}(X_2) = 4$ that $\dim \text{End}(X_{22}) + \dim \text{Hom}(X_{21}, X_{22}) \leq 3$.

On the other hand, by Proposition 4.1.2, we have

$$0 \longrightarrow \tau^{-1}P(1) \oplus X_{21} \longrightarrow E_1 \bigoplus \oplus 2E_2 \longrightarrow X_{22} \longrightarrow 0.$$

Applying $\text{Hom}(\cdot, X_{22})$ (resp. $\text{Hom}(E_1 \bigoplus \oplus 2E_2, \cdot)$) to the exact sequence above, we get $\dim \text{Hom}_\Lambda(E_1 \bigoplus \oplus 2E_2, X_{22}) \leq 4$ (resp. $\dim \text{Hom}(E_1 \bigoplus \oplus 2E_2, X_{22}) \geq 5$) by $\text{Hom}(\tau^{-1}P(1), X_2) = k$ (resp. $\text{Hom}(E_1 \bigoplus \oplus 2E_2, \tau^{-1}P(1) \oplus X_{21}) = 0$).

It implies that X_2 contains a summand X'_2 with $X'_2 \in \mathcal{T}_1$.

Assume that $X_2 = E_1 + X'_2$, then there does not exist any pre-projective direct summand M of X'_2 such that $\text{Hom}(M, P) = 0$ and $\exists f \in \text{Hom}(P, M)$ with f to be an epimorphism by the AR -quiver of $\tilde{E}_6$.

If X'_2 does not contain a summand of regular module. It follows from Proposition 4.1.2 that

$$0 \longrightarrow \tau^{-1}P(1) \longrightarrow \oplus 2E_2 \longrightarrow X'_2 \longrightarrow 0.$$

Applying $\text{Hom}(\cdot, \oplus 2E_2)$, we get $4 = \dim \text{End}(\oplus 2E_2) \leq 2 = \dim \text{Hom}(\tau^{-1}P(1), \oplus 2E_2)$. Therefore, $X \cong \tau^{-1}P(1) \oplus E_1 \oplus E_2 \oplus \tau I(1)$.

Assume that $X_2 = E_2 + X'_2$. When X'_2 contains a pre-projective summand M such that $\text{Hom}(M, P) = 0$ and $\exists f \in \text{Hom}(P, M)$ with f to be an epimorphism, we can deduce $M \cong \tau^{-3}P(1)$ by the AR -quiver of $\tilde{E}_6$. Thus $X \cong \tau^{-1}P(1) \oplus E_2 \oplus \tau^{-3}P(1) \oplus I(5) \oplus I(7)$ and $\dim \text{End}(X) \geq 7$, which gives a contradiction.

When X'_2 has only pre-injective summands, we get

$$0 \longrightarrow \tau^{-1}P(1) \longrightarrow E_1 \oplus E_2 \longrightarrow X'_2 \longrightarrow 0.$$

Applying $\text{Hom}(\cdot, E_1 \oplus E_2)$ to the exact sequence above, we get a contradiction $2 = \dim \text{End}(E_1 \oplus E_2) \leq 1 = \dim \text{Hom}(\tau^{-1}P(1), E_1 \oplus E_2)$.

From the discussions above, we get $X \cong \tau^{-1}P(1) \oplus E_1 \oplus E_2 \oplus \tau I(1)$.

Thus

$$\begin{aligned} p_3|_{\overline{\mathcal{O}}_{E_1 \oplus \oplus 2E_2} \times \mathcal{O}_S}^{-1}(\tau^{-1}P(1) \oplus \tau^{-3}P(2)) &= V(S, E_1 \bigoplus \oplus 2E_2; \tau^{-1}P(1) \oplus \tau^{-3}P(2)) \\ &\cup V(S, \tau I(5) \oplus \tau I(7); \tau^{-1}P(1) \cup (p_3|_{\overline{\mathcal{O}}_{E_1 \oplus \oplus E_2} \times \mathcal{O}_{P(2)}})^{-1}(\tau^{-3}P(2))). \end{aligned}$$

By $m = r = 1$, we deduce that $(p_3|_{\overline{\mathcal{O}}_{E_1 \oplus E_2} \times \mathcal{O}_{P(2)}})^{-1}(\tau^{-3}P(2))$ has purity property. Thus $p_3|_{\overline{\mathcal{O}}_{E_1 \oplus \oplus 2E_2} \times \mathcal{O}_S}^{-1}(\tau^{-1}P(1) \oplus \tau^{-3}P(2))$ has also the purity property.

Suppose $P = P_1 \oplus P_2 \oplus P_3$. Without loss of generality, let $P = \tau^{-1}P(1) \oplus \tau^{-2}P(1) \oplus \tau^{-3}P(1)$. Because S_1 is a summand of $\text{top}(E_1/\text{Im}(f))$ for any $f \in \text{Hom}(P_1 \oplus P_2 \oplus P_3, E_1)$, we obtain that $P_1 \oplus P_2 \oplus P_3$ is not an extension of S by $E_1 \bigoplus \oplus 2E_2$. The proof for Case 3 is complete.

Case 4 : $Q = \tilde{E}_7$.

We only consider $\mathcal{T}_2$ in Fig 4.2.4. The other cases can be proved similarly. Let $E'_i, i = 1, 2, 3$, be simple objects in the full subcategory $\mathcal{T}_2$ of $\Lambda - \text{mod}$ corresponding to dimension vectors listed in Fig 4.2.4.

Since

$$\dim_k \text{Ext}^1(E'_1, S) = \dim_k \text{Ext}^1(E'_2, S) = 1, \dim_k \text{Ext}^1(E'_3, S) = 2,$$

we have

$$\dim_k \text{Ext}^1(\oplus m E'_1, S) = \dim_k \text{Ext}^1(\oplus m E'_2, S) = m, \dim_k \text{Ext}^1(\oplus m E'_3, S) = 2m.$$

In addition, we have

$$\begin{aligned} \theta : \quad 0 &\longrightarrow S \xrightarrow{\begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}} P(1) \oplus P(7) \oplus P(8) \longrightarrow E'_1 \longrightarrow 0, \\ \theta' : \quad 0 &\longrightarrow S \xrightarrow{\begin{bmatrix} 1 \\ -1 \end{bmatrix}} P(2) \oplus P(6) \longrightarrow E'_2 \longrightarrow 0 \end{aligned}$$

be a basis of $\text{Ext}^1(E'_1, S)$ and $\text{Ext}^1(E'_2, S)$, respectively.

Obviously, $P(1) \oplus P(7) \oplus P(8) \oplus \oplus (m-1)E'_1$ (resp. $P(2) \oplus P(6) \oplus \oplus (m-1)E'_2$) is a non-trivial extension of S by $\oplus m E'_1$ (resp. $\oplus m E'_2$).

Based on the claim above, we may assume that $P \oplus (\oplus (m-a)E'_1 \oplus \oplus (p-b)E'_2 \oplus \oplus (r-c)E'_3)$ is a non-trivial extension of S by $\oplus m E'_1 \oplus \oplus p E'_2 \oplus \oplus r E'_3$, for some a, b, c , where P is pre-projective.

Up to isomorphism, $(P(1) \oplus P(7) \oplus P(8)) \oplus \oplus (m-1)E'_1$ (resp. $P(2) \oplus P(6) \oplus \oplus (m-1)E'_2$) is the unique non-trivial extension of S by $\oplus m E'_1$ (resp. $\oplus m E'_2$).

Since

$$p_3^{-1}|_{\overline{\mathcal{O}}_{\oplus m E'_1} \times \mathcal{O}_S} (P(1) \oplus P(7) \oplus P(8)) = P^{2m}, p_3|_{\overline{\mathcal{O}}_{\oplus m E'_2} \times \mathcal{O}_S}^{-1} (P(2) \oplus P(6)) = P^m, ,$$

we deduce they have purity property.

Because E'_3 is non-sincere, we can reduce the study of $\text{Ext}_{\Lambda}^1(\oplus m E'_3, S)$ to the study of $\text{Ext}_{\Lambda'}^1(\oplus m E'_3, S)$, where Λ' is the path algebra of Dynkin quiver D_4 determined by vertexes 3, 4, 5 and 8.

By case 3, we get

$$\begin{cases} \mathbf{Z}_{\oplus m E'_3, S, X_1 \oplus \oplus (m-1)E'_3} = (\mathbb{A}^{m+1} - \cup_{j=1}^3 \mathcal{L}(r_{N_j}))/k^*, \\ \mathbf{Z}_{\oplus m E'_3, S, (P(8) \oplus Y_1) \oplus \oplus (m-1)E'_3} = (\mathbb{A}^{m+1} - \cup_{j=1}^3 \mathcal{L}(r_{N_j}))/k^*, \end{cases}$$

and

$$\begin{cases} p_3^{-1}|_{\overline{\mathcal{O}}_{\oplus m E'_3} \times \mathcal{O}_S} (X_1 \oplus \oplus (m-1)E'_3) \\ \quad = P^{\dim_k \text{Hom}(S, X_1 \oplus \oplus (m-1)E'_3) - 1}, \\ p_3^{-1}|_{\overline{\mathcal{O}}_{\oplus m E'_3} \times \mathcal{O}_S} (P(8) \oplus Y_1 \oplus \oplus (m-1)E'_3) \\ \quad = P^{\dim_k \text{Hom}(S, P(8) \oplus Y_1 \oplus \oplus (m-1)E'_3) - 1} \end{cases}$$

Thus $p_3^{-1}|_{\overline{\mathcal{O}}_{\oplus m E'_3} \times \mathcal{O}_S} (X_1 \oplus \oplus (m-1)E'_3)$ and $p_3^{-1}|_{\overline{\mathcal{O}}_{\oplus m E'_3} \times \mathcal{O}_S} (P(8) \oplus Y_1 \oplus \oplus (m-1)E'_3)$ have purity property.

Similarly, the statement is true if $mp \neq 0, r = 0$ or $pr \neq 0, m = 0$ or $rm \neq 0, p = 0$.

We now assume that $mpr \neq 0$. Because of (A) and Mayer-Vietoris sequence, we only need to consider $a = m, b = p, c = r$.

Consider the short exact sequence

$$(5.15) \quad 0 \longrightarrow S \longrightarrow P \longrightarrow \oplus mE'_1 \oplus \oplus pE'_2 \oplus \oplus rE'_3 \longrightarrow 0.$$

Applying $H(\oplus mE'_1 \oplus \oplus pE'_2 \oplus \oplus rE'_3, \cdot)$, $i = 1, 2, 3$ to (5.15), we get

$$m^2 + p^2 + r^2 \leq m + p + 2r.$$

Thus $m = p = 1, 1 \leq r \leq 2$.

When $m = p = r = 1$.

Consider the short exact sequence

$$0 \longrightarrow S \longrightarrow P \longrightarrow E'_1 \oplus E'_2 \oplus E'_3 \longrightarrow 0.$$

Because $\langle \delta, \dim P \rangle = -4$, we know that P contains 4 indecomposable objects at most.

Suppose that P is indecomposable. Let $X \in \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3}$ such that P is an extension of S by X . Thus we have the following exact sequence

$$(5.16) \quad 0 \longrightarrow S \longrightarrow P \longrightarrow X \longrightarrow 0.$$

Applying $\text{Hom}(\cdot, S)$ (resp. $\text{Hom}(X, \cdot)$) to (5.16), we get from $\text{Hom}(P, S) = 0$ and $\text{Hom}(X, P) = 0$ that

$$\text{Hom}(X, S) = 0, \text{Ext}^1(X, S) = k^4, \dim \text{End}(X) = 4.$$

If $X \cong X_1 \oplus X_2$ with X_i indecomposable for all $i = 1, 2$, we have $X_1 \in \mathcal{P}_{\text{prep}}, X_2 \in \mathcal{P}_{\text{prei}}$ and $\text{Hom}(X_1, X_2) = 2$. Thus $X_1 \cong \tau^{-3}P(3)$ or $\tau^{-3}P(5)$, and $X \cong \tau^{-3}P(3) \oplus I(3)$ or $\tau^{-3}P(5) \oplus I(5)$.

Without loss of generality, we may assume that $X \cong \tau^{-3}P(3) \oplus I(3)$.

Fix one exact sequence

$$0 \longrightarrow P(1) \longrightarrow P \longrightarrow \tau^{-3}P(3) \longrightarrow 0.$$

Because $\dim \text{Hom}(P(1), I(3)) = 1, \dim \text{Hom}(S, P(1)) = 1$, in the same way as (I) in case 3, we get

$$V(S, \tau^{-3}P(3) \oplus I(3); P) \cong V(S, I(3); P(1)) = \{1 \text{ point}\}.$$

If $X \cong X_1 \oplus X_2 \oplus X_3$ with X_i indecomposable for all $i = 1, 2, 3$. Assume that there is not regular module in X_i . Without loss of generality, let $X_1, X_2 \in \mathcal{P}_{\text{prep}}$ and $X_3 \in \mathcal{P}_{\text{prei}}$. Because of Proposition 4.1.2 and $\dim \text{End}(X) = 4$, we have

$$0 \longrightarrow X_1 \oplus X_2 \longrightarrow E'_1 \oplus E'_2 \oplus E'_3 \longrightarrow X_3 \longrightarrow 0.$$

Applying $\text{Hom}(\cdot, X_3)$ (resp. $\text{Hom}(E'_1 \oplus E'_2 \oplus E'_3, \cdot)$) to the sequence above, we get

$$\begin{aligned} \dim_k \text{Hom}_\Lambda(E'_1 \oplus E'_2 \oplus E'_3, X_3) &\leq 2, \\ \dim_k \text{Hom}_\Lambda(E'_1 \oplus E'_2 \oplus E'_3, X_3) &\geq 3. \end{aligned}$$

This give a contradiction. Thus $X_1 \in \mathcal{P}_{\text{prep}}, X_2 \in \mathcal{T}_2, X_3 \in \mathcal{P}_{\text{prei}}$.

When $X_2 \cong E'_1$. Because of $\text{Hom}(X, P) = 0$, it follows from the AR -quiver of $\tilde{E}_7$ that $X_1 \cong \tau^{-5}P(1)$ or $X_1 \cong \tau^{-5}P(7)$. Thus

$$X \cong \tau^{-5}P(1) \oplus E'_1 \oplus \tau^2I(1) \text{ or } X \cong \tau^{-5}P(7) \oplus E'_1 \oplus \tau^2I(7).$$

By $\text{End}(\tau^{-5}P(1) \oplus E'_1 \oplus \tau^2I(1)) = \text{End}(\tau^{-5}P(7) \oplus E'_1 \oplus \tau^2I(7)) = k^3$, This give contradiction. When $X_2 \cong E'_2$. We get that X_1 is isomorphic to $\tau^{-3}P(8)$ or $\tau^{-i}P(j)$ for $i = 3, 4, 5, 7; j = 1, 7$.

Suppose $X_1 \cong \tau^{-3}P(8)$, and if P is the extension of S by $\tau^{-3}P(8) \oplus E'_2 \oplus I(8)$. Since

$$\dim_k \text{Hom}_\Lambda(P, \tau^{-3}P(8)) = \dim_k \text{Hom}_\Lambda(P, E'_2) = 1, \dim_k \text{Hom}_\Lambda(P, I(8)) = 2,$$

$$\text{Aut}_\Lambda(E'_1 \oplus E'_2 \oplus E'_3) = \left\{ \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} \mid a \in \text{Aut}(E'_1), b \in \text{Aut}(E'_2), c \in \text{Aut}(E'_3) \right\}$$

we get

$$V(S, \tau^{-3}P(8) \oplus E'_2 \oplus I(8); P) = \mathbb{A}^1 \setminus V_0, |V_0| < \infty, \text{ or } V(S, \tau^{-3}P(8) \oplus E'_2 \oplus I(8); P) = \emptyset.$$

Suppose $X_1 \cong \tau^{-3}P(1)$ or $X_1 \cong \tau^{-3}P(7)$, then we know that X has 4 indecomposable summands at least.

Suppose $X_1 \cong \tau^{-4}P(1)$ or $X_1 \cong \tau^{-4}P(7)$, then we have $\text{End}(X) = 3$.

Suppose $X_1 \cong \tau^{-5}P(1)$ or $X_1 \cong \tau^{-5}P(7)$, then P has the direct summands $I(1)$ or $I(7)$.

Suppose $X_1 \cong \tau^{-7}P(1)$ or $X_1 \cong \tau^{-7}P(7)$, we then have $X \cong \tau^{-7}P(1) \oplus E'_2 \oplus I(1)$ or $X \cong \tau^{-7}P(7) \oplus E'_2 \oplus I(7)$. Since $\text{Ext}_\Lambda^1(I(1), S) = \text{Ext}_\Lambda^1(I(7), S) = 0$, we have P to be decomposable.

Therefore, P is only possibly the extension of S by $\tau^{-3}P(8) \oplus E'_2 \oplus I(8)$ in this case.

When $X_2 \cong E'_3$. We get

$$X \cong \tau^{-3}P(2) \oplus E'_3 \oplus I(2), \text{ or } \tau^{-3}P(1) \oplus E'_3 \oplus \tau^4 I(1), \text{ or } \tau^{-6}P(1) \oplus E'_3 \oplus \tau I(1),$$

$$X \cong \tau^{-3}P(6) \oplus E'_3 \oplus I(6), \text{ or } \tau^{-3}P(7) \oplus E'_3 \oplus \tau^4 I(7), \text{ or } \tau^{-6}P(7) \oplus E'_3 \oplus \tau I(7).$$

Suppose $X \cong \tau^{-3}P(2) \oplus E'_3 \oplus I(2)$, or $\tau^{-3}P(6) \oplus E'_3 \oplus I(6)$.

Because of $\text{Ext}^1(I(2), S) = \text{Ext}^1(I(6), S) = 0$, then $I(2)$ or $I(6)$ is the direct summands of P .

Suppose

$$X \cong \tau^{-3}P(1) \oplus E'_3 \oplus \tau^4 I(1), \text{ or } \tau^{-3}P(7) \oplus E'_3 \oplus \tau^4 I(7),$$

then we get $\text{End}(X) = 3$.

Suppose

$$X \cong \tau^{-6}P(1) \oplus E'_3 \oplus \tau I(1), \text{ or } \tau^{-6}P(7) \oplus E'_3 \oplus \tau I(7).$$

It follows from $\text{Ext}^1(\tau I(1), S) = \text{Ext}^1(\tau I(7), S) = 0$ that P is a decomposable.

Thus P is also not an extension of S by $X_1 \oplus E'_3 \oplus X_3$.

If $X \cong X_1 \oplus X_2 \oplus X_3 \oplus X_4$ with X_i indecomposable for all $i = 1, 2, 3, 4$, then we have that $\{X_4, X_3, X_2, X_1\}$ is an orthogonal exceptional sequence. Thus $\mathfrak{C}(X_4, X_3, X_2, X_1)$ is isomorphic to $k\tilde{A}_3 - \text{mod}$. Because of $X \in \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3}$ and $\dim \text{End}(X) = 4$, we see that $\{X_1, X_2, X_3, X_4\}$ are simple objects in $\mathfrak{C}(X_4, X_3, X_2, X_1)$, and $X_1 \cong \tau^{-3}P(7)$, $X_2 \cong E'_2$, $X_3 \cong E'_3$, $X_4 \cong I(7)$ or $X_1 \cong \tau^{-3}P(1)$, $X_2 \cong E'_2$, $X_3 \cong E'_3$, $X_4 \cong I(1)$. Therefore P must be decomposable. This also give a contradiction.

From above, we get

$$(p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3} \times \mathcal{O}_S})^{-1}(P)$$

$$= V(S, E'_1 \oplus E'_2 \oplus E'_3; P) \cup V(S, \tau^{-3}P(3) \oplus I(3); P)$$

$$\cup V(S, \tau^{-3}P(5) \oplus I(5); P) \cup V(S, \tau^{-3}P(8) \oplus E'_2 \oplus I(8); P).$$

Since

$$V(S, E'_1 \oplus E'_2 \oplus E'_3; P) = \mathbb{A}^1 \setminus V_0, |V_0| < \infty,$$

we deduce that $(p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3} \times \mathcal{O}_S})^{-1}(P)$ has purity property.

Suppose $P \cong P_1 \oplus P_2$ with P_i indecomposable. Without loss of generality, we assume that $P \cong P(7) \oplus \tau^{-3}P(5)$.

Let $X \in \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3} \setminus \mathcal{O}_{E'_1 \oplus E'_2 \oplus E'_3}$ such that $P(7) \oplus \tau^{-3}P(5)$ is an extension of S by X .

If $\text{Hom}(X, P(7) \oplus \tau^{-3}P(5)) = 0$, similar to the case of P indecomposable, we can deduce that $P(7) \oplus \tau^{-3}P(5)$ is not an extension of S by X .

If $\text{Hom}(X, P(7) \oplus \tau^{-3}P(5)) \neq 0$, we then get $X \cong P(7) \oplus X_2$ or $X \cong \tau^{-3}P(5) \oplus I(5)$ by the AR -quiver of $\tilde{E}_7$.

When $X \cong P(7) \oplus X_2$, it is easy to see that $V(S, X; P(7) \oplus \tau^{-3}P(5)) = V(S, X_2; \tau^{-3}P(5))$.

When $X \cong \tau^{-3}P(5) \oplus I(5)$, we get $V(S, X; P(7) \oplus \tau^{-3}P(5)) = V(S, I(5); P(7)) = \{1 \text{ point}\}$.

Thus

$$\begin{aligned} & (p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3} \times \mathcal{O}_S})^{-1}(P(7) \oplus \tau^{-3}P(5)) \\ &= V(S, E'_1 \oplus E'_2 \oplus E'_3; P(7) \oplus \tau^{-3}P(5)) \cup_{[X_2]} (p_3|_{\overline{\mathcal{O}}_{X_2} \times \mathcal{O}_S})^{-1}(\tau^{-3}P(5)) \cup V(S, I(5); P(7)). \end{aligned}$$

By [L4], we know that $(p_3|_{\overline{\mathcal{O}}_{X_2} \times \mathcal{O}_S})^{-1}(\tau^{-3}P(5))$ has purity property. Since

$$V(S, E'_1 \oplus E'_2 \oplus E'_3, P(7) \oplus \tau^{-3}P(5)) = \mathbb{A}^1 \setminus V_0, |V_0| < \infty,$$

we can deduce that $(p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3} \times \mathcal{O}_S})^{-1}(P(7) \oplus \tau^{-3}P(5))$ has purity property.

Suppose that $P \cong P_1 \oplus P_2 \oplus P_3$ with P_i indecomposable for all $i = 1, 2, 3$. Without loss of generality, we assume that $P \cong P(7) \oplus \tau^{-1}P(7) \oplus \tau^{-3}P(6)$.

Let $X \in \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3} \setminus \mathcal{O}_{E'_1 \oplus E'_2 \oplus E'_3}$ such that $P(7) \oplus \tau^{-1}P(7) \oplus \tau^{-3}P(6)$ is an extension of S by X .

If $\text{Hom}(X, P(7) \oplus \tau^{-1}P(7) \oplus \tau^{-3}P(6)) = 0$, similar to the case of P indecomposable, we can deduce that $P(7) \oplus \tau^{-1}P(7) \oplus \tau^{-3}P(6)$ is not an extension of S by X .

If $\text{Hom}(X, P(7) \oplus \tau^{-1}P(7) \oplus \tau^{-3}P(6)) \neq 0$, we then know that X contains a direct summand $P(7)$, or $\tau^{-1}P(7)$, or $\tau^{-2}(P(7))$, or $\tau^{-2}(P(6))$, or $\tau^{-3}(P(6))$.

When $X \cong \tau^{-2}(P(7)) \oplus X_2$ with X_2 indecomposable, we have

$$\text{Hom}(\tau^{-2}(P(7)), X_2) = \text{Hom}((P(7)), \tau^2 X_2) = 0.$$

Thus $\dim \text{End}(X) = 2$, this give a contradiction.

When $X \cong \tau^{-2}(P(7)) \oplus X_2 \oplus X_3$ with X_2, X_3 to be indecomposable modules, we then have $X_2 \in \mathcal{T}_2$.

Suppose $X \cong \tau^{-2}(P(7)) \oplus E'_1 \oplus X_3$. Then $\text{End}(X) = 3$, it forces $\tau^{-2}(P(7)) \oplus E'_1 \oplus X_3 \notin \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3} \setminus \mathcal{O}_{E'_1 \oplus E'_2 \oplus E'_3}$.

Suppose $X \cong \tau^{-2}(P(7)) \oplus E'_2 \oplus X_3$ (resp. $X \cong \tau^{-2}(P(7)) \oplus E'_3 \oplus X_3$.) Then $I(1)$ (resp. $\tau I(7)$) is a direct summands of X_3 . It implies $I(1)$ (resp. $\tau I(7)$) to be a direct summands of P . This also contradicts to the assumption.

Similarly, we may show that P is not a extension of S by X if $\tau^{-1}P(7)$ or $\tau^{-2}P(6)$ or $\tau^{-3}P(6)$ is a direct summands of X .

Since

$$V(S, E'_1 \oplus E'_2 \oplus E'_3; P_1 \oplus P_2 \oplus P_3) = \mathbb{A}^1 \setminus V_0, |V_0| < \infty,$$

we get

$$(p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3} \times \mathcal{O}_S})^{-1}(P_1 \oplus P_2 \oplus P_3) = V(S, E'_1 \oplus E'_2 \oplus E'_3; P_1 \oplus P_2 \oplus P_3)$$

has purity property.

Suppose $P \cong P_1 \oplus P_2 \oplus P_3 \oplus P_4$ with P_i indecomposable for all $i = 1, 2, 3, 4$. Without loss of generality, we may assume that $P \cong P(7) \oplus \tau^{-1}P(7) \oplus \tau^{-2}P(7) \oplus \tau^{-3}P(7)$.

Because $\text{Hom}(P(7) \oplus \tau^{-1}P(7) \oplus \tau^{-3}P(7), E'_2) = 0$ and there is not an epimorphism from $\tau^{-2}P(7)$ to E'_2 , it implies that $P(7) \oplus \tau^{-1}P(7) \oplus \tau^{-2}P(7) \oplus \tau^{-3}P(7)$ is not an extension of S by $E'_1 \oplus E'_2 \oplus E'_3$.

When $m = p = 1, r = 2$, let P be the extension of S by $E'_1 \oplus E'_2 \oplus 2E'_3$. Because of $\langle \delta, \underline{\dim} S \rangle = \langle \delta, \underline{\dim} P \rangle = -4$, we can deduce that P has 4 indecomposable direct summands at most.

Assume that P is also the extension of S by X with $X \in \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus 2E'_3} \setminus \mathcal{O}_{E'_1 \oplus E'_2 \oplus 2E'_3}$. Then there is an exact sequence

$$(5.17) \quad 0 \longrightarrow S \longrightarrow P \longrightarrow X \longrightarrow 0.$$

Because of $\text{Hom}(P, S) = 0$, we deduce that $\text{Hom}(X, S) = 0, \text{Ext}^1(X, S) = k^6$. Applying $\text{Hom}(X, \cdot)$ to (5.17), we get $\dim \text{End}(X) \leq 6 + \dim \text{Hom}(X, P)$.

Suppose P is an indecomposable. We then have $\dim \text{End}(X) \leq 6$. Thus $X \notin \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus 2E'_3}$. This give a contradiction.

In the same computing as in case 3, we have

$$(p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus 2E'_3} \times \mathcal{O}_S})^{-1}(P) = V(S, E'_1 \oplus E'_2 \oplus 2E'_3; P)$$

has purity property.

Suppose $P \cong P_1 \oplus P_2$ with P_i indecomposable. Without loss of generality, we may set $P \cong \tau^{-1}P(7) \oplus \tau^{-4}P(5)$.

If $\text{Hom}(X, P) = 0$, it follows that $\dim \text{End}(X) \leq 6$ and $X \notin \overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus 2E'_3}$. This give a contradiction.

Thus $\text{Hom}(X, P) \neq 0$, and we get $\tau^{-4}P(5) \vdash X$ or $\tau^{-1}P(7) \vdash X$.

When $X \cong \tau^{-4}P(5) \oplus X'$, we deduce that $X \cong I(8) \oplus \tau^2 I(1) \oplus \tau^{-4}P(5)$. Thus

$$V(S, I(8) \oplus \tau^2 I(1) \oplus \tau^{-4}P(5); \tau^{-1}P(7) \oplus \tau^{-4}P(5)) = V(S, I(8) \oplus \tau^2 I(1); \tau^{-1}P(7)) = \{1 \text{ point}\}.$$

When $X \cong \tau^{-1}P(7) \oplus X_2$. We get $\dim \text{Hom}(X, P) = 1$ and $\dim \text{End}(X) = 7$. We now set $X \cong \tau^{-1}P(7) \oplus X_2$. It follows from $\text{Hom}(\tau^{-1}P(7), X_2) = 1$ that $\text{End}(X_2) = 5$ and X_2 is decomposable.

Assume that $X_2 \cong X_{21} \oplus X_{22}$ with $X_{21} \in \mathcal{P}_{\text{prep}}, X_{22} \in \mathcal{P}_{\text{prei}}$. By $\text{End}(X_2) = 5$, we get $\dim \text{End}(X_{22}) + \dim \text{Hom}(X_{21}, X_{22}) \leq 4$. Based on Proposition 4.1.2, we have

$$0 \longrightarrow \tau^{-1}P(7) \oplus X_{21} \longrightarrow E'_1 \oplus E'_2 \oplus 2E'_3 \longrightarrow X_{22} \longrightarrow 0.$$

Applying $\text{Hom}(\cdot, X_{22})$ (resp. $\text{Hom}(E'_1 \oplus E'_2 \oplus 2E'_3, \cdot)$) to the sequence above, we get

$$\dim \text{Hom}(E'_1 \oplus E'_2 \oplus 2E'_3, X_{22}) \leq 5 \text{ and } \dim \text{Hom}(E'_1 \oplus E'_2 \oplus 2E'_3, X_{22}) \geq 6.$$

This give a contradiction. Thus X_2 must contain one regular direct summand at least.

According to the AR -quiver of $\tilde{E}_7$, we can deduce that $X \cong \tau^{-1}P(7) \oplus E'_1 \oplus E'_2 \oplus E'_3 \oplus \tau^2 I(7)$.

Thus

$$\begin{aligned} p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3 \oplus \tau^2 I(7)} \times \mathcal{O}_S}^{-1}(\tau^{-1}P(7) \oplus \tau^{-4}P(5)) &= V(S, E'_1 \oplus E'_2 \oplus E'_3 \oplus \tau^2 I(7); \tau^{-1}P(7) \oplus \tau^{-4}P(5)) \\ &\cup V(S, I(8) \oplus \tau^2 I(1); \tau^{-1}P(7)) \cup (p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3} \times \mathcal{O}_{P(5)}})^{-1}(\tau^{-4}P(5)). \end{aligned}$$

By $m = p = r = 1$, we deduce that $(p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3} \times \mathcal{O}_{P(5)}})^{-1}(\tau^{-4}P(5))$ has purity property. So does $p_3|_{\overline{\mathcal{O}}_{E'_1 \oplus E'_2 \oplus E'_3 \oplus \tau^2 I(7)} \times \mathcal{O}_S}^{-1}(\tau^{-1}P(7) \oplus \tau^{-4}P(5))$.

Suppose $P \cong P_1 \oplus P_2 \oplus P_3$ with P_i indecomposable. Without loss of generality, we may set $P \cong \tau^{-1}P(7) \oplus \tau^{-2}P(7) \oplus \tau^{-4}P(6)$.

For any $f \in \text{Hom}(\tau^{-2}P(7), E'_2)$, we have $S(6) \vdash E'_2 / \text{Im}(f)$.

By $\text{Hom}(\tau^{-1}P(7) \oplus \tau^{-4}P(6), E'_2) = 0$, we deduce that $\tau^{-1}P(7) \oplus \tau^{-2}P(7) \oplus \tau^{-4}P(6)$ is not an extension of S by $E'_1 \oplus E'_2 \oplus \oplus 2E'_3$.

Similarly, suppose $P \cong P_1 \oplus P_2 \oplus P_3 \oplus P_4$ with P_i indecomposable, then we may show that there is not an epimorphism from P to $E'_1 \oplus E'_2 \oplus \oplus 2E'_3$. Thus P is not an extension of S by $E'_1 \oplus E'_2 \oplus \oplus 2E'_3$.

The proof is complete. $\square$

Remark 5.3 Let P be non-trivial extension of $\oplus tS$ by the regular semi-simple objects in $\mathcal{T}_i$ for some i .

Since

$$\begin{aligned} \mathbf{Z}_{S, \oplus(t-1)S; \oplus tS} &= \mathbb{P}^1, \\ \mathbb{P}^1 \times (p_3|_{\overline{\mathcal{O}}_M * \mathcal{O}_{\oplus tS}})^{-1}(P) \\ &= \cup_{P' \oplus L, L \vdash M} (p_3|_{\overline{\mathcal{O}}_M * \mathcal{O}_S})^{-1}(P' \oplus L) \times (p_3|_{\overline{\mathcal{O}}_{P' \oplus L} * \mathcal{O}_{\oplus(t-1)S}})^{-1}(P) \\ &\quad \cup (p_3|_{\overline{\mathcal{O}}_{P'' \oplus L' \oplus I', P'' \oplus L' \oplus I' \in \overline{\mathcal{O}}_M, L' \vdash M} * \mathcal{O}_{\oplus tS}})^{-1}(P). \end{aligned}$$

By induction on t and $\dim M$, Proposition 5.2 is true when S is replaced by $\oplus mS$. $\square$

Proposition 5.4 Let i be a sink, and P, P' pre-projective modules of kQ (except $Q = \tilde{E}_8$), and let M be a regular semi-simple modules in $\mathcal{T}_i$ for some $i, 1 \leq i \leq l$.

Then $p_3|_{\overline{\mathcal{O}}_M * \mathcal{O}_P}^{-1}(P' \oplus L)$ have purity property, where L is a submodule of M in $\mathcal{T}_i$.

Proof. Let $P = \oplus a_1 P_1 \oplus \oplus a_2 P_2 \oplus \oplus \cdots \oplus \oplus a_t P_t$ with $\text{Ext}(P_i, P_j) = 0$ for $i < j$ and P_i is indecomposable.

First, suppose that $P = \oplus a_1 P_1$. Applying reflection functor σ_i , we can turn the question into that of in case $P = \oplus a_1 P(i)$. By Proposition 5.2 and remark 5.3, the statement is true.

By the properties of pre-projective components, the statement above is true if M is replaced $P' \oplus M$, where P' is pre-projective.

Next, suppose that $P \cong \oplus a_1 P_1 \oplus \oplus a_2 P_2 \oplus \oplus \cdots \oplus \oplus a_t P_t$ and $t \geq 2$. Without loss generality, we may only consider the case $t = 2$.

Since

$$p_3|_{\overline{\mathcal{O}}_M * \mathcal{O}_{\oplus a_1 P_1 \oplus \oplus a_2 P_2}}^{-1}(P' \oplus L) = \cup_{P'' \oplus L', L' \vdash M} (p_3|_{\overline{\mathcal{O}}_M * \mathcal{O}_{a_1 P_1}})^{-1}(P'' \oplus L') \times (p_3|_{\overline{\mathcal{O}}_{P'' \oplus L} * \mathcal{O}_{a_2 P_2}})^{-1}(P' \oplus L),$$

it follows from case $t = 1$ that $p_3|_{\overline{\mathcal{O}}_M * \mathcal{O}_{\oplus a_1 P_1 \oplus \oplus a_2 P_2}}^{-1}(P' \oplus L)$ have purity property. The proof is complete. $\square$

Dually, we also have the following statement

Proposition 5.5 Let i be a source, and I, I' pre-injective modules of kQ (except $Q = \tilde{E}_8$), and let M be a regular semi-simple modules in $\mathcal{T}_i$ for some $i, 1 \leq i \leq l$.

Then $p_3|_{\overline{\mathcal{O}}_M * \mathcal{O}_I}^{-1}(I' \oplus L)$ have purity property, where L is a submodule of M in $\mathcal{T}_i$. $\square$

Proposition 5.6 Let i be a sink, and P, P' pre-projective modules of kQ (except $Q = \tilde{E}_8$), and let M be a regular module in $\mathcal{T}_i$ for some $i, 1 \leq i \leq l$.

Then $p_3|_{\overline{\mathcal{O}}_M * \mathcal{O}_P}^{-1}(P' \oplus L)$ have purity property, where L is a submodule of M in $\mathcal{T}_i$.

Proof. By using induction on $\dim M$, we prove that $p_3|_{\overline{\mathcal{O}}_M * \mathcal{O}_P}^{-1}(P' \oplus L)$ have purity property.

The case that M is semi-simple object in $\mathcal{T}_i$ is proved in Proposition 5.4.

Assume that M is not semi-simple in $\mathcal{T}_i$. Let M' be the direct sum of indecomposable summands of M with maximal length in full subcategory $\mathcal{T}_i$ of $\Lambda - \text{mod}$. Set $M_2 = \text{soc}_{\mathcal{T}_i}(M')$.

According to Proposition 2.5 in [GJ], there is a regular module M_1 in $\mathcal{T}_i$ such that $g_{M_1 M_2}^M = 1$.

Since

$$\begin{cases} \mathcal{O}_{M_1} * \mathcal{O}_{M_2} = p_3 p_2 p_1^{-1}(\mathcal{O}_{M_1} \times \mathcal{O}_{M_2}), \\ \mathbf{Z}_{M, M_1, M_2} = p_2 p_1^{-1}(\mathcal{O}_{M_1} \times \mathcal{O}_{M_2}) \cap p_3^{-1}(M) = \{1 \text{ point}\}, \end{cases}$$

and $\mathcal{O}_{M_1} * \mathcal{O}_{M_2}$ has only finitely many orbits, we thus get

$$(5.18) \quad \begin{cases} \dim \mathcal{O}_{M_1} * \mathcal{O}_{M_2} = \dim \mathcal{O}_M, \\ \overline{\mathcal{O}}_{M_1} * \overline{\mathcal{O}}_{M_2} = \overline{\mathcal{O}}_M. \end{cases}$$

Thus

$$\begin{aligned} p_3|_{\overline{\mathcal{O}}_M * \mathcal{O}_P}^{-1}(P' \oplus L) &= p_3|_{\overline{\mathcal{O}}_{M_1} * \overline{\mathcal{O}}_{M_2} * \mathcal{O}_P}^{-1}(P' \oplus L) \\ &= \cup_{P'' \oplus L', L' \vdash M_2} p_3|_{\overline{\mathcal{O}}_{M_1} * \mathcal{O}_{P'' \oplus L'}}^{-1}(P' \oplus L) \times p_3|_{\overline{\mathcal{O}}_{M_2} * \mathcal{O}_P}^{-1}(P'' \oplus L'). \end{aligned}$$

Since M_2 is semi-simple in full subcategory, the disjoint union above make sense. By Proposition 5.4, we have that $p_3|_{\overline{\mathcal{O}}_{M_2} * \mathcal{O}_P}^{-1}(P'' \oplus L')$ have purity property.

In addition, it follows from the induction hypothesis and the proof of proposition 2.5 in [GJ] that $p_3|_{\overline{\mathcal{O}}_{M_1} * \mathcal{O}_{P'' \oplus L'}}^{-1}(P' \oplus L)$ have purity property.

Thus the proof is complete. $\square$

6. Canonical bases of $\mathcal{H}^s(\Lambda)$

6.1 This subsection is offered to prove 4.1.5.

Lemma 6.1.1 *Let M (resp. N) be a module belong to the nonhomogeneous tubes $\mathcal{T}_i$ (resp. $\mathcal{T}_j$) in $kQ - \text{mod}$ (except $Q = \tilde{E}_8$) for $i \neq j$, and $\overline{\mathcal{O}}_M$ (resp. $\overline{\mathcal{O}}_N$) have purity property.*

Then $\overline{\mathcal{O}}_{M \oplus N}$ have purity property.

Proof. Set $\alpha = \underline{\dim} M, \beta = \underline{\dim} N$, and $X = \overline{\mathcal{O}}_{M \oplus N}$. Let

$$P = \left\{ \begin{bmatrix} A & C \\ 0 & B \end{bmatrix} \middle| \begin{bmatrix} A & C \\ 0 & B \end{bmatrix} \in GL_{\alpha+\beta}, A \in GL_{\alpha}, B \in GL_{\beta} \right\}.$$

Then P is a parabolic subgroup of $GL_{\alpha+\beta}$.

It is well known that the closure Y of $GL_{\alpha+\beta}/P$ has purity property.

Consider the natural projection

$$p : GL_{\alpha+\beta}/(Aut_{\Lambda}(M) \times Aut_{\Lambda}(N)) \longrightarrow GL_{\alpha+\beta}/P.$$

It is easy to see that the fibre of p is $P/(Aut(M) \times Aut(N))$.

By the definition of $Aut_{\Lambda}(M)$ and $Aut_{\Lambda}(N)$, we have

$$\begin{aligned} Aut_{\Lambda}(M) &\subseteq GL_{\alpha}, \quad Aut_{\Lambda}(N) \subseteq GL_{\beta}, \\ P/(Aut_{\Lambda}(M) \times Aut_{\Lambda}(N)) &\cong GL_{\alpha}/Aut_{\Lambda}(M) \times GL_{\beta}/Aut_{\Lambda}(N) \times \prod_{i \in I} k^{\alpha_i \beta_i}. \end{aligned}$$

Because the closure of $GL_{\alpha}/Aut_{\Lambda}(M)$ and $GL_{\beta}/Aut_{\Lambda}(N)$ have purity property, we can deduce that the closure Z of $P/(Aut_{\Lambda}(M) \times Aut_{\Lambda}(N))$ also have purity property.

Applying it to the long exact sequence

$$\cdots \longrightarrow H_c^i(Z, \overline{\mathbb{Q}}_l) \longrightarrow H_c^i(X, \overline{\mathbb{Q}}_l) \longrightarrow H_c^i(Y, \overline{\mathbb{Q}}_l) \longrightarrow \cdots,$$

the proof is complete. $\square$

Note that $Hom_{\Lambda}(M, N)$ is a subspace of $\prod_{i \in I} Hom_k(k^{\alpha_i}, k^{\beta_i})$. In the same way as in Lemma 5.4, we have

Lemma 6.1.2 *Let M and N be modules of kQ (except $Q = \tilde{E}_8$), $Ext_{\Lambda}^1(M, N) = 0$, $Hom_{\Lambda}(N, M) = 0$ and $\overline{\mathcal{O}}_M$ (resp. $\overline{\mathcal{O}}_N$) have purity property. Then $\overline{\mathcal{O}}_{M \oplus N}$ have purity property. $\square$*

Proposition 6.1.3 *Let M be a regular kQ - (except $Q = \tilde{E}_8$) module in $\mathcal{T}_i$ for $1 \leq i \leq l$. Then $\overline{\mathcal{O}}_M$ have purity property.*

Proof. We use induction on $\dim M$.

The case that of M is semi-simple object in $\mathcal{T}_i$ is proved in Lemma 4.2.1-Lemma 4.2.5.

Assume that M is not semi-simple in $\mathcal{T}_i$. Let M' be the direct sum of indecomposable summands of M with maximal length in full subcategory $\mathcal{T}_i$ of $\Lambda - \text{mod}$. Set $M_2 = \text{soc}_{\mathcal{T}_i}(M')$.

According to Proposition 2.5 in [GJ], there is a regular module M_1 in $\mathcal{T}_i$ such that $g_{M_1 M_2}^M = 1$. It follows from (5.18) that $\overline{\mathcal{O}}_{M_1} * \overline{\mathcal{O}}_{M_2} = \overline{\mathcal{O}}_M$.

By induction hypothesis and Lemma 4.2.1-Lemma 4.2.5, we obtain $\overline{\mathcal{O}}_{M_1}$ and $\overline{\mathcal{O}}_{M_2}$ have purity property.

On the other hand, because there are finitely many orbits in $\overline{\mathcal{O}}_{M_2}$, for any $P \oplus R \oplus I \in \overline{\mathcal{O}}_{M_2}$, by proposition 5.6 and the proof of Proposition 2.5 in [GJ], $(p_3|_{\overline{\mathcal{O}}_{M_1} * \mathcal{O}_{P \oplus R}})^{-1}(P' \oplus R')$ has purity property. So does $(p_3|_{\overline{\mathcal{O}}_{M_1} * \mathcal{O}_{P \oplus R \oplus I}})^{-1}(P' \oplus R' \oplus I')$.

Thus

$$(p_3|_{\overline{\mathcal{O}}_{M_1} * \overline{\mathcal{O}}_{M_2}})^{-1}(P' \oplus R' \oplus I') = \cup_{P \oplus R \oplus I \in \overline{\mathcal{O}}_{M_2}} (p_3|_{\overline{\mathcal{O}}_{M_1} * \mathcal{O}_{P \oplus R \oplus I}})^{-1}(P' \oplus R' \oplus I').$$

Consider the proper morphism

$$p_3 : p_2 p_1^{-1}(\overline{\mathcal{O}}_{M_1} \times \overline{\mathcal{O}}_{M_2}) \longrightarrow \overline{\mathcal{O}}_{M_1} * \overline{\mathcal{O}}_{M_2}.$$

From the discussions above, we obtain $\overline{\mathcal{O}}_M$ has purity property.

We now prove Theorem 4.1.5. By [L4] and [L5], $\overline{\mathcal{O}}_P$ (resp. $\overline{\mathcal{N}}_{\mathbf{w},3}$, $\overline{\mathcal{O}}_I$) have purity property. It now follows from Proposition 6.1.3 that $\overline{\mathcal{O}}_M$ have purity property.

According to Lemma 6.1.2, $X = \overline{\mathcal{O}}_{P,M,\mathbf{w},I}$ have purity property, that is, the desired conclusion follows from above. $\square$

6.2 Let

$$(6.2.1) \quad b_{\overline{\mathcal{O}}_{P,M,\mathbf{w},I}} = \sum_{i, N \in \overline{\mathcal{O}}_{P,M,\mathbf{w},I}^F} (-1)^i v^{i + \dim \mathcal{O}_N} \dim \mathcal{H}^i(IC_{\overline{\mathcal{O}}_{P,M,\mathbf{w},I}}(\overline{\mathbb{Q}}_l)) \langle N \rangle.$$

Set

$$\mathbf{CB} = \{b_{\overline{\mathcal{O}}_{P,M,\mathbf{w},I}} | P \in \mathcal{P}_{\text{prep}}, M \in \oplus_{i=1}^l \mathcal{T}_i, I \in \mathcal{P}_{\text{prei}}, \mathbf{w} \in \mathbf{P}(n), n \in \mathbb{N}\}.$$

Then we have the main theorem in this paper

Proposition 6.2.1 *The set $\mathbf{CB}$ is the canonical basis of $\mathcal{H}^s(\Lambda)$ (except $Q = \tilde{E}_8$).*

Proof. By Proposition 1.3.1, Theorem 3.1.1 and Proposition 6.1.3, the proof is complete. $\square$

7. Application of $\mathcal{H}^s(\Lambda)$

7.1 In [H], A. Hubery have proved the existence of Hall polynomials on the tame quivers for Segre classes. In this subsection, by using the extension algebras of singularity Ringel-Hall algebras, we give a simple and direct proof for the existence of Hall polynomials on the tame quivers. .

Let Q be a quiver. We define a new algebra $\mathcal{H}^{s,x_1,x_2,\dots,x_t}$ which is generated by $\{u_i, u_{[M]}, u_{[N]} : i \in I, M \in \mathcal{T}_j, N \in \oplus_t \mathcal{H}_{x_t}, 1 \leq j \leq l, t \in \mathbb{N}\}$, where x_t are $\mathbb{F}_q$ rational points in $\mathbb{P}^1$.

We now give a new decomposition of $E_{n\delta}$ as follows

$$E_{n\delta} = E_{n\delta,1} + E_{n\delta,2} + E_{n\delta,3},$$

where

$$E_{n\delta,1} = v^{-n \dim S_1 - n \dim S_2} \sum_{[M], M \in \mathfrak{C}_1 \oplus \oplus_{x_t} \mathcal{H}_{x_t}, \underline{\dim} M = n\delta} u[M], \quad (7.1.1)$$

$$E_{n\delta,2} = v^{-n \dim S_1 - n \dim S_2} \sum_{\substack{[M], \underline{\dim} M = n\delta, \\ M = M_1 \oplus M_2, 0 \neq M_1 \in \mathfrak{C}_1 \oplus \oplus_{x_t} \mathcal{H}_{x_t}, 0 \neq M_2 \in \mathfrak{C}_0 \setminus \oplus_{x_t} \mathcal{H}_{x_t}}} u[M], \quad (7.1.2)$$

$$E_{n\delta,3} = v^{-n \dim S_1 - n \dim S_2} \sum_{[M], M \in \mathfrak{C}_0 \setminus \oplus_{x_t} \mathcal{H}_{x_t}, \underline{\dim} M = n\delta} u[M]. \quad (7.1.3)$$

Let $\mathbf{w} = (w_1, \dots, w_t)$ be a partition of n , we then define

$$E_{\mathbf{w}\delta,3} = E_{w_1\delta,3} * \dots * E_{w_t\delta,3}.$$

Let $\mathbf{P}(n)$ be the set of all partitions of n , and $\langle N \rangle = v^{-\dim N + \dim \text{End}(N)} u_{[N]}$. Set

$$\mathbf{B}' = \{ \langle P \rangle * \langle M \rangle * E_{\mathbf{w}\delta,3} * \langle I \rangle \mid P \in \mathcal{P}_{\text{prep}}, M \in \oplus_{i=1}^l \mathcal{T}_i \bigoplus \oplus_{x_t} \mathcal{H}_{x_t}, I \in \mathcal{P}_{\text{prei}}, \mathbf{w} \in \mathbf{P}(n), n \in \mathbb{N} \}.$$

Similar to Theorem 3.1.1, we have the following:

Proposition 7.1.1 *The set $\mathbf{B}'$ is $\mathbb{Q}(v)$ -basis of $\mathcal{H}^{s,x_1,x_2,\dots,x_t}$.* $\square$

Theorem 7.1.2 *Let Q be a quiver, let P_i (resp. R_i, I_i) be a pre-projective (resp. nonhomogeneous regular, pre-injective) $\mathbb{F}_q Q$ -modules, and let $H_i \in \oplus_{j=1}^t \mathcal{H}_{x_j}$ be homogeneous regular $\mathbb{F}_q Q$ -modules with x_i to be $\mathbb{F}_q$ -rational points in $\mathbb{P}^1$ for $i = 1, 2, 3; j \in \mathbb{N}$.*

Then there exists Hall polynomial $\varphi_{12}^3(x) \in \mathbb{Q}[x]$ such that

$$\varphi_{12}^3(q) = g_{P_1 \oplus R_1 \oplus H_1 \oplus I_1, P_2 \oplus R_2 \oplus H_2 \oplus I_2}^{P_3 \oplus R_3 \oplus H_3 \oplus I_3}.$$

Proof. Since

$$\langle P_1 \oplus M_1 \oplus I_1 \rangle * \langle P_2 \oplus M_2 \oplus I_2 \rangle = a_{12}^3(v) \langle P_3 \oplus M_3 \oplus I_3 \rangle + \text{the other terms},$$

and $a_{12}^3(v) \in \mathbb{Q}(v)$. Then we have

$$\begin{aligned} & g_{P_1 \oplus R_1 \oplus H_1 \oplus I_1, P_2 \oplus R_2 \oplus H_2 \oplus I_2}^{P_3 \oplus R_3 \oplus H_3 \oplus I_3} \\ &= v^{\dim_{\mathbb{F}_q} \text{End}(P_3 \oplus M_3 \oplus I_3) - \dim_{\mathbb{F}_q} \text{End}(P_1 \oplus M_1 \oplus I_1) - \dim_{\mathbb{F}_q} \text{End}(P_2 \oplus M_2 \oplus I_2) - \langle \underline{\dim} P_1 \oplus R_1 \oplus H_1 \oplus I_1, \underline{\dim} P_2 \oplus R_2 \oplus H_2 \oplus I_2 \rangle} a_{12}^3(v). \end{aligned}$$

Set

$$(*) \quad \begin{aligned} & \varphi_{12}^3(v^2) \\ &= v^{\dim_{\mathbb{F}_q} \text{End}(P_3 \oplus M_3 \oplus I_3) - \dim_{\mathbb{F}_q} \text{End}(P_1 \oplus M_1 \oplus I_1) - \dim_{\mathbb{F}_q} \text{End}(P_2 \oplus M_2 \oplus I_2) - \langle \underline{\dim} P_1 \oplus R_1 \oplus H_1 \oplus I_1, \underline{\dim} P_2 \oplus R_2 \oplus H_2 \oplus I_2 \rangle} a_{12}^3(v). \end{aligned}$$

On the other hand, we known that

$$v^{\dim_{\mathbb{F}_q} \text{End}(P_3 \oplus M_3 \oplus I_3) - \dim_{\mathbb{F}_q} \text{End}(P_1 \oplus M_1 \oplus I_1) - \dim_{\mathbb{F}_q} \text{End}(P_2 \oplus M_2 \oplus I_2) - \langle \underline{\dim} P_1 \oplus R_1 \oplus H_1 \oplus I_1, \underline{\dim} P_2 \oplus R_2 \oplus H_2 \oplus I_2 \rangle} a_{12}^3(v)$$

takes the positive integer value while v^2 takes infinite many positive integer values. Then $\varphi_{12}^3(v^2)$ is the polynomial of v^2 over $\mathbb{Q}$. Thus the proof is complete. $\square$

REFERENCES

- [BBD] A. Beilinson, I.N. Bernstein, P.D. Deligne *Faisceaux Pervers*, Astérisque **100** (1981).
- [BN] J. Beck and H. Nakajima, *Crystal bases and two sided cells of quantum affine algebras*, Duke Math. J. **123** (2004), 335-402.
- [BGP] I. N. Bernstein, I. M. Gelfand and V. A. Ponomarev, *Coxeter functors and Gabriel theorem*, Uspechi Mat. Nauk **28** (1973); Russian Math. Surv. **28** (1973), 17–32.
- [CB] W. Crawley-Boevey, *Lectures on Representations of Quivers*, Mathematical Institute, Oxford University, 1992
- [DDX] B. Deng, J. Du and J. Xiao, *Generic extensions and canonical bases for cyclic quivers*, to appear in Can. J. Math.
- [DR] V. Dlab and C. M. Ringel, *Indecomposable representations of graphs and algebras*, Memoirs Amer. Math. Soc. **173**, 1976.
- [G] J. A. Green, *Hall algebras, hereditary algebras and quantum groups*, Invent. Math. **120** (1995), 361–377.
- [H] G.J. A. Hubery, *Hall polynomials for affine quivers*, arXiv:math/0703178v2.
- [GJ] J. Y. Guo, *The Hall polynomials of a cyclic serial algebra*, Comm. Alg. **23**(2) (1995), 743–751.
- [HX] J. Hua and J. Xiao, *On Ringel-Hall algebra of tame hereditary algebra*, Algebr. Represent. Theory **5**(2002),no.5,527-550.
- [K] M. Kashiwara, *On crystal bases of the q -analogue of universal enveloping algebras*, Duke Math. J. **63**(1991), 465–516.
- [KW] R. Kiehl and R. Weissauer, *Weil conjectures, Perverse sheaves and l' adic Fourier transform*, A Series of Modern surveys in Mathematics, Vol **42**, 2000.
- [LXZ] Z. Lin, J. Xiao and G. Zhang *Representations of tame quivers and affine canonical bases*, Preprint, arXiv:math.QA/0706.1444v3.
- [L1] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), 447–498.
- [L2] G. Lusztig, *Quivers, perverse sheaves, and the quantized enveloping algebras*, J. Amer. Math. Soc. **4** (1991), 366–421.
- [L3] G. Lusztig, *Affine quivers and canonical bases*, Inst. Hautes Études Sci. Publ. Math. **76** (1992), 111–163.
- [L4] G. Lusztig, *Canonical bases and Hall algebras*, A. Broer and A. Daigneault(eds.), Representation Theories and Algebraic Geometry, 365–399, 1998 Kluwer Academic Publishers.
- [L5] G. Lusztig, *Introduction to Quantum Groups*, Birkhäuser, Boston, 1993.
- [N] H. Nakajima, *Crystal, canonical and PBW bases of quantum affine algebras*, Preprint.
- [Re] M. Reineke, *The monoid of families of quiver representations*, Proc. London, Math. Soc., **3(84)** (2002), 663–685.
- [R1] C. M. Ringel, *Hall algebras and quantum groups*, Invent. math. **101** (1990), 583–592.
- [R2] C. M. Ringel, *Hall algebras*, In: Topics in Algebra, Banach Center Publ. **26** (1990), 433–447.
- [R3] C. M. Ringel, *Hall polynomials for representation finite hereditary algebras*, Adv. Math. **84** (1990), 137–178.
- [R4] C. M. Ringel, *The preprojective algebra of a tame quiver: The irreducible components of the module varieties*, Contemporary Math. Vol **229** (1990), 293–306.
- [SV] B. Sevenhant and M. Van den Bergh, *A relation between a conjecture of Kac and the structure of the Hall algebra*, J. Pure and Appl. Algebra **160** (2001), 319–332.
- [VV] M. Varagnolo and E. Vasserot, *On the decomposition matrices of the quantized Schur algebra*, Duke Math. J. **100** (1999), 267–297.
- [X] J. Xiao, *Drinfeld double and RingelGreen theory of Hall algebras*, J. Algebra **190**(1997), 100–144.
- [Z] Z. Zwara, *Degenerations of finite dimensional modules are given by extensions*, compos. Math. **121**(2000), 205-218.

DEPARTMENT OF MATHEMATICS, SHANGHAI JIAOTONG UNIVERSITY, SHANGHAI 200240, CHINA

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, TAIPEI, TAIWAN 11529

E-mail address: g.l.zhang@sjtu.edu.cn